



# OFFICIAL PRODUCTS

BROCHURE 2024



# WHO ARE WE?



## OUR MESSAGE

We provide high quality feed additives, vitamins with the best price, and as fast as possible, all over the world. ADEMIX is a private limited liability company, established in 2010 in Mersin, Turkey with the efforts and support of experts and veterinary specialists. ADEMIX provides veterinary products in Turkey and all over the world, and seeks to be a leader in this field.



## COMPANY STRATEGY

Establishing mutual cooperation relations with our clients, exchanging technical and scientific experiences, and contributing to preserving human health through the development of animal health sector.





## COMPANY VALUES

Our Company Values is:



- Human health through animal health.



- the success of our customers is our success.



## OUR VISION

We aim to be a world leader in securing materials for the veterinary industry with rapid demand security, high material quality and competitive prices.



## OUR MISSION

We provide feed additives with the highest quality, and best price, and as fast as possible, all over the world.



# 2023 CALENDAR

## January

S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

## February

S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28				

## March

S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

## April

S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30						

## May

S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31			

## June

S	M	T	W	T	F	S
			1	2	3	
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	

## July

S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

## August

S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

## September

S	M	T	W	T	F	S
				1	2	
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30

## October

S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				

## November

S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30		

## December

S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					





# 2024 CALENDAR

## January

S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31			

## February

S	M	T	W	T	F	S
				1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29		

## March

S	M	T	W	T	F	S
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30
31						

## April

S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30				

## May

S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

## June

S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30						

## July

S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31			

## August

S	M	T	W	T	F	S
				1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31

## September

S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30					

## October

S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

## November

S	M	T	W	T	F	S
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30

## December

S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30	31				



# LIQUIDS PRODUCTS

ADECK-VIT

VITAMIX - 100

VITAMIX - 50

VITAMIX - 25

VITAMIX - 10

VITAMIX - 5

SELEN E - 20 %

SELEN E MIX

MINOPLEX

AMINOVIT MAX

VITAMINO FORT

MAXI POWER

SHELLMIN D3

ADEPHOS

ADEMIN SOL

CARVA MENT

OREGO MIX - 5

OKASAK

IMMUNE STAR

IMMUNEMAX

LIVOMIX

LIVOMIX FORT

HEPAMIX

DIUREMIX

BITKI POWER

BREATHMIX

ELECTROSOL

ANTIKET

ADETOX - S

ACID MIX

ANTI MOLD



1000 ml

# ADECK-VIT



## COMPOSITIONS

Each 1000 ml contains:

Vitamin A	40,000,000	IU
Vitamin D3	5,000,000	IU
Vitamin E	50,000	mg
Vitamin C	25,000	mg
Vitamin K	25,000	mg
Excipients to	1,000	ml

phosphorus, which are required for normal growth, bone development, and eggshell formation.

- Vitamin E is a powerful antioxidant and important for normal neurological functions.

Deficiency leads to encephalomalacia/crazy chick disease, muscular dystrophy seen more frequently in older and mature birds.

- Vitamin C ameliorates heat stress induced problems such as poor immunity, feed intake, weight gain, oxidative stress, rectal and body temperature, fertility and semen quality, carcass weight and mortality in birds.

- Some of the vitamin K is synthesized by intestinal bacteria. It is necessary for blood coagulation. It helps prevent osteoporosis and bone fractures. Its deficiency can cause bleeding disorders and anemia.



## BENEFITS:

- Vitamin A is required for normal growth, reproduction and maintenance of epithelial cells in good condition (skin and the linings of the digestive, reproductive, and respiratory tracts).

- Vitamin D3 is required for proper absorption and utilization of calcium and

## IT'S USED FOR:



## INDICATIONS:

- It is used in cases of vitamin deficiency.
- Keeping poultry in good health.
- Increasing production.
- Raising fertility.
- As an aid to the formation of immunity.



## DOSAGE:

- Poultry: 500 ml per 1000 liters of drinking water from 3-5 day.
- Cattle and Horses: 10 ml per 100 Kg bodyweight for 3-5 day.
- Sheep, Goats and Calves: 10 ml per 100 Kg bodyweight for 3-5 day.







1000 ml

#### IT'S USED FOR:



# VITAMIX-100



## COMPOSITIONS

Each 1000 ml contains:

Vitamin A	100,000,000	IU
Vitamin D3	20,000,000	IU
Vitamin E	25,000	mg
Excipients to	1,000	ml

- Vitamin A is required for normal growth, reproduction and maintenance of epithelial cells in good condition (skin and the linings of the digestive, reproductive, and respiratory tracts).
- Vitamin D3 is required for proper absorption and utilization of calcium and phosphorus, which are required for normal growth, bone development, and eggshell formation.
- Vitamin E is a powerful antioxidant and important for normal neurological functions. Deficiency leads to encephalomalacia/crazy chick disease, muscular dystrophy seen more frequently in older and mature birds.



## INDICATIONS:

- It is used in cases of vitamin deficiency.
- Keeping ruminants in good health.
- Keeping poultry in good health.
- Increasing production.
- Raising fertility.
- As an aid to the formation of immunity.



## BENEFITS:

- Vitamin A is required for normal growth, reproduction and maintenance of epithelial cells in good condition (skin and the linings of the digestive, reproductive, and respiratory tracts).



## DOSAGE:

- Poultry: 150 ml per 1000 liters of drinking water from 3-5 day.
- Horses and Cattle: 100-150 ml per 1000 liters of drinking water from 3-5 day.
- Sheep, Goats and Calves: 10 ml per 1000 liters of drinking water for 3-5 days.







1000 ml

#### IT'S USED FOR:



# VITAMIX - 50



## COMPOSITIONS

Each 1000 ml contains:

Vitamin A	50,000,000	IU
Vitamin D3	2,500,000	IU
Vitamin E	25,000	mg
Excipients to	1,000	ml

- Vitamin A is required for normal growth, reproduction and maintenance of epithelial cells in good condition (skin and the linings of the digestive, reproductive, and respiratory tracts).
- Vitamin D3 is required for proper absorption and utilization of calcium and phosphorus, which are required for normal growth, bone development, and eggshell formation.
- Vitamin E is a powerful antioxidant and important for normal neurological functions. Deficiency leads to encephalomalacia/crazy chick disease, muscular dystrophy seen more frequently in older and mature birds.



## INDICATIONS:

- It is used in cases of vitamin deficiency.
- Keeping poultry in good health.
- Increasing production.
- Raising fertility.
- As an aid to the formation of immunity.



## BENEFITS:

- Vitamin A is required for normal growth, reproduction and maintenance of epithelial cells in good condition (skin and the linings of the digestive, reproductive, and respiratory tracts).
  - Vitamin D3 is required for proper absorption and utilization of calcium and phosphorus, which are required for normal growth, bone development, and eggshell formation.
  - Vitamin E is a powerful antioxidant and important for normal neurological functions. Deficiency leads to encephalomalacia/crazy chick disease, muscular dystrophy seen more frequently in older and mature birds.
- Poultry: 500 ml per 1000 liters of drinking water from 3-5 day.
- Cattle and Horses: 10 ml per 100 Kg bodyweight for 3-5 day.
- Ruminants: 1000 ml per 1000 liters of drinking water from 3-5 day.



## DOSAGE:



# VITAMIX - 25



## COMPOSITIONS

Each 1000 ml contains:

Vitamin A	25,000,000	IU
Vitamin D3	7,000,000	IU
Vitamin E	25,000	mg
Excipients to	1,000	ml

- Vitamin D3 is required for proper absorption and utilization of calcium and phosphorus, which are required for normal growth, bone development, and eggshell formation.
- Vitamin E is a powerful antioxidant and important for normal neurological functions deficiency leads to encephalomalacia/crazy chick disease, muscular dystrophy seen more frequently in older and mature birds.



## INDICATIONS:

- It is used in cases of vitamin deficiency.
- Keeping ruminants in good health.
- Keeping poultry in good health.
- Increasing production.
- Raising fertility.
- As an aid to the formation of immunity.



## BENEFITS:

- Vitamin A is required for normal growth, reproduction and maintenance of epithelial cells in good condition (skin and the linings of the digestive, reproductive, and respiratory tracts).



## DOSAGE:

- Poultry: 1000 ml per 1000 liters of drinking water from 3-5 day.
- Cattle and Horses: 5 - 10 ml per animal during 3 - 5 day.
- Sheep, Goats and Calves: 2 - 3 ml per animal during 3 - 5 day.

## IT'S USED FOR:







1000 ml

#### IT'S USED FOR:



# VITAMIX - 10



#### COMPOSITIONS

Each 1000 ml contains:

Vitamin A	10,000,000	IU
Vitamin D3	5,000,000	IU
Vitamin E	5,000	mg
Excipients to	1,000	ml

- Vitamin D3 is required for proper absorption and utilization of calcium and phosphorus, which are required for normal growth, bone development, and eggshell formation.
- Vitamin E is a powerful antioxidant and important for normal neurological functions deficiency leads to encephalomalacia/crazy chick disease, muscular dystrophy seen more frequently in older and mature birds.



#### INDICATIONS:

- It is used in cases of vitamin deficiency.
- Keeping ruminants in good health.
- Keeping poultry in good health.
- Increasing production.
- Raising fertility.
- As an aid to the formation of immunity.



#### BENEFITS:

- Vitamin A is required for normal growth, reproduction and maintenance of epithelial cells in good condition (skin and the linings of the digestive, reproductive, and respiratory tracts).



#### DOSAGE:

- Poultry: 1500 ml per 1000 liters of drinking water from 3-5 day.
- Cattle and Horses: 10 - 15 ml per animal during 3 - 5 day.
- Sheep, Goats and Calves: 3 - 5 ml per animal during 3 - 5 day.





# VITAMIX - 5



## COMPOSITIONS

Each 1000 ml contains:

Vitamin A	5,000,000	IU
Vitamin D3	1,000,000	IU
Vitamin E	5,000	mg
Excipients to	1,000	ml

- Vitamin D3 is required for proper absorption and utilization of calcium and phosphorus, which are required for normal growth, bone development, and eggshell formation.
- Vitamin E is a powerful antioxidant and important for normal neurological functions deficiency leads to encephalomalacia/crazy chick disease, muscular dystrophy seen more frequently in older and mature birds.



## INDICATIONS:

- It is used in cases of vitamin deficiency.
- Keeping ruminants in good health.
- Keeping poultry in good health.
- Increasing production.
- Raising fertility.
- As an aid to the formation of immunity.



## BENEFITS:

- Vitamin A is required for normal growth, reproduction and maintenance of epithelial cells in good condition (skin and the linings of the digestive, reproductive, and respiratory tracts).



## DOSAGE:

- Poultry: 2000 ml per 1000 liters of drinking water from 3-5 day.
- Cattle and Horses: 15 - 20 ml per animal during 3 - 5 day.
- Sheep, Goats and Calves: 5 - 10 ml per animal during 3 - 5 day.

## IT'S USED FOR:







# SELEN E - 20%



## COMPOSITIONS

Each 1000 ml contains:

Vitamin E	200,000	mg
Selenium	900	mg
Excipients to	1,000	ml



## SYMPTOMS OF A VITAMIN E DEFICIENCY INCLUDE:

Crazy chick disease (encephalomalacia), Wry neck, Decreased laying Muscular dysfunction, Weakness, Inability to walk, bowed legs, Greenish -blue skin, Muscle spasms, Inability to control head (to turn it, hold it up or lower it) Hemorrhages, Paralysis, Enlarged hocks, Poor growth, Poor feed conversion Increase susceptibility to disease and parasites and Decreased fertility.



## BENEFITS:

- Vitamin E is a powerful antioxidant and important for normal neurological functions, Greater resistance to heat stress, strengthened immune response, including to E. coli, Improved laying, A better hatch rates, Higher levels of vitamin E in eggs and meat.
- Selenium has been shown to have a strong antioxidant, anti-inflammatory, and antimicrobial ability.



## INDICATIONS:

- It is used in cases of vitamin E deficiency in calves, lambs, sheep, goats, piglets and poultry.
- Encephalomalacia (Crazy Chick Disease).
- Muscular dystrophy (White Muscle Disease, Stiff Lamb Disease).
- Decreased hatchability of eggs.
- Dietary selenium supplementation has been used for improving growth performance, immune function, and meat quality.
- Selenium is an essential trace mineral important for the maintenance of homeostasis in animals.



## DOSAGE:

- Poultry: 500 ml /1000 liters of drinking water from 3-5 day.
- Calves: 1 – 3 ml / 1 liters for 3-5 day.
- Cows: 7.5 - 15 ml / 1 liters for 3-5 day.
- Sheep: 2 - 5 ml / 1 liters for 3-5 day.
- Lambs: 1 – 3 ml / 1 liters for 3-5 day.

## IT'S USED FOR:





# SELEN E MIX



## COMPOSITIONS

Each 1000 ml contains:

Vitamin E	100,000	mg
Selenium	450	mg
Excipients to	1,000	ml



## SYMPTOMS OF

a vitamin E deficiency include:

Crazy chick disease (encephalomalacia), Wry neck, Decreased laying Muscular dysfunction, Weakness, Inability to walk, bowed legs, Greenish -blue skin, Muscle spasms, Inability to control head (to turn it, hold it up or lower it) Hemorrhages, Paralysis, Enlarged hocks, Poor growth, Poor feed conversion Increase susceptibility to disease and parasites and Decreased fertility.



## INDICATIONS:

- It is used in cases of vitamin E deficiency in calves, lambs, sheep, goats, piglets and poultry.
- Encephalomalacia (Crazy Chick Disease).
- Muscular dystrophy (White Muscle Disease, Stiff Lamb Disease).
- Decreased hatchability of eggs.
- Dietary selenium supplementation has been used for improving growth performance, immune function, and meat quality.
- Selenium is an essential trace mineral important for the maintenance of homeostasis in animals.



## BENEFITS:

- Vitamin E is a powerful antioxidant and important for normal neurological functions, Greater resistance to heat stress, strengthened immune response, including to E. coli, Improved laying, A better hatch rates, Higher levels of vitamin E in eggs and meat.
- Selenium has been shown to have a strong antioxidant, anti-inflammatory, and antimicrobial ability.



## DOSAGE:

- Poultry: 1000 ml /1000 liters of drinking water from 3-5 day.
- Calves: 3 - 5 ml / 1 liters for 3-5 day.
- Cows: 20 - 25 ml / 1 liters for 3-5 day.
- Sheep: 5 - 10 ml / 1 liters for 3-5 day.
- Lambs: 2 - 5 ml / 1 liters for 3-5 day.

## IT'S USED FOR:





# MINOPLEX



## COMPOSITIONS

Each 1000 ml contains:

Vitamin B2	200	mg
Niacin	3,500	mg
Calcium-D-Pantothenate	130	mg
Vitamin B6	125	mg
Vitamin B12	1,250	mcg
DL-Methionine	1,000	mg
L-Lysine Hcl	1,000	mg
Choline Chloride 75%	1,000	mg
Excipients to	1,000	ml



## BENEFITS:

- The B vitamins are involved in many metabolic functions, including energy metabolism, a vitamin premix is typically used to compensate for the fluctuating levels of vitamins found naturally in food and to assure adequate levels of all vitamins.
- Riboflavin (Vitamin B2) is part of enzyme systems so plays a vital role in metabolism.
- Pyridoxine (Vitamin B6) is necessary for proper metabolism of amino acids.

- Cyanocobalamin (Vitamin B12) is involved with nucleic acid synthesis, carbohydrate and fat metabolism and methyl synthesis.
- Methionine and lysine: is an essential amino acids as it cannot be synthesized by the bird, it serves as a constituent of body protein and is involved in the development of the digestive tract and growth performance.

## IT'S USED FOR:



## INDICATIONS:

- Promoting appetite.
- Weight gain.
- Increased productivity.



## DOSAGE:

- Poultry:  
1000 ml / 1000 liters of drinking water for 3-5 day.
- Calves and Lambs:  
20 ml / 100 kg body weight.







# AMINOVIT MAX



## COMPOSITIONS

Each 1000 ml contains:

Vitamin A	14,000,000	IU	L-Threonine	490	mg
Vitamin D3	600,000	IU	L-Valin	920	mg
Vitamin E	10,000	mg	L-Alanine	650	mg
Vitamin B1	3,000	mg	L-Arginine	810	mg
Vitamin B2	500	mg	Aspartic Acid	700	mg
Niacin	10,000	mg	L-Phenylalanine	600	mg
Calcium- D-Pantothenate	4,000	mg	Glycine	1,030	mg
Vitamin B6	1,000	mg	Monosodium Glutamate	1,410	mg
Biotin	15	mg	L-Isoleucine	540	mg
Vitamin B12	20	mg	L-Leucine	970	mg
Vitamin C	7,000	mg	L-Lysine Hcl	1,730	mg
Choline Chloride	8,000	mg	L-Proline	1,240	mg
Carnitine	1,500	mg	L-Tyrosine	324	mg
DL-Methionine	378	mg	Excipients to	1,000	ml

## IT'S USED FOR:



## INDICATIONS:

• AMINOVIT MAX is used as a supplementary source of essential amino acids, fat-soluble vitamins such as A-D3-E, and water-soluble vitamins such as B-vitamin complexes and vitamin C, in poultry, horses, calves and lambs.



## DOSAGE:

- Poultry: 500 ml / 1000 liters of drinking water for 3 - 5 days.
- Calves and Lambs: 10 ml / 100 kg body weight daily.
- Horses: 10 ml / 100 kg body weight 3 - 5 days before the race.







# VITAMINO FORT



## COMPOSITIONS

Each 1000 ml contains:

Vitamin A	20,000,000	IU	L-Threonine	122.5	mg
Vitamin D3	2,000,000	IU	Monosodium Glutamate	500	mg
Vitamin E	8,000	mg	L-Proline	310	mg
Vitamin B1	3,000	mg	L-Glycine	257.5	mg
Vitamin B2	2,400	mg	L-Leucine	242.5	mg
Niacin	12,000	mg	L-Valin	230	mg
Calcium- D-Pantothenate	6,000	mg	L-Arginine	202.5	mg
Vitamin B6	1,200	mg	L-Aspartic Acid	175	mg
Vitamin B12	6,000	mcg	L-Alanine	162.5	mg
Vitamin C	10,000	mg	L-Phenylalanine	150	mg
Magnesium Chloride	1,000	mg	L-Isoleucine	135	mg
Zinc Oxide	313	mg	L-Tyrosine	81	mg
Sodium Iodate	6	mg	Yeast Extract	50,000	mg
Iron Chloride	294	mg	Propylene Glycol	20,000	mg
L-Lysine Hcl	432.5	mg	Excipients to	1,000	ml
DL-Methionine	94.5	mg			

## IT'S USED FOR:



## INDICATIONS:

It is used as a nutritional supplement to meet the animal's needs for vitamins, amino acids and trace minerals.



## DOSAGE:

- Poultry: 500 ml / 1000 liters of drinking water for 3 – 5 days.
- Calves and Lambs: 10 ml / 100 kg body weight daily.
- Horses: 10 ml / 100 kg body weight 3-5 days before the race.





1000 ml

# MAXI POWER



## COMPOSITIONS

Each 1000 ml contains:

Vitamin A	8,000,000	IU	Potassium	600	mg
Vitamin D3	1,000,000	IU	Sodium	1,180	mg
Vitamin E	10,000	mg	Magnesium	3,200	mg
Vitamin B1	4,000	mg	Beta Glucan	4,000	mg
Vitamin B2	1,000	mg	Mannan Oligosaccharides	4,000	mg
Niacin	6,000	mg	DL-Methionine	945	mg
Choline Chloride	6,500	mg	L-Threonine	1,225	mg
Calcium- D-Pantothenate	4,000	mg	L-Valin	2,300	mg
Vitamin B6	1,000	mg	L-Alanine	1,625	mg
Biotin	15	mg	L-Arginine	2,025	mg
Vitamin B12	5	mg	Aspartic Acid	1,750	mg
Vitamin C	900	mg	L-Phenylalanine	1,500	mg
Carnitine	1,200	mg	L-Glycine	2,575	mg
Iron	1,800	mg	Monosodium Glutamate	3,525	mg
Zinc	4,000	mg	L-Isoleucine	1,360	mg
Copper	100	mg	L-Lucien	2,425	mg
Manganese	230	mg	L-Lysine Hcl	4,325	mg
Phosphor	2,400	mg	L-Proline	3,100	mg
Cobalt	50	mg	L-Thyrosine	810	mg
Iodine	100	mg	Aroma	10,000	mg
Selenium	200	mg	Excipients to	1,000	ml

FULL OF



• AMINO ACIDS • VITAMINS  
• MINERALS ★ LIVER SUPPORT

IT'S USED FOR:







## INDICATIONS:

It is used as a Supplementary Ground in order to meet the vitamin, amino acid, trace mineral and aromatic substance needs of Adult Rams, Sheep, Goats, Tosun, Dairy Cows and Beef Cattles, and Lambs, Capricorns and Calves that have completed their rumen development.



## DOSAGE:

- Adult Rams and Goats 10 - 12 ml per animal per day.
- Adult Sheep and Goats 6 - 7 ml per animal per day.
- 3 - 4 ml per animal per day for Lambs and Kids that have completed their rumen development.
- Adult Tosun Dairy Cows and Fattening Cattles 10 - 12 ml per 100 kg body weight per day.
- It is used by mixing 6 - 7 ml of drinking water per animal per day for calves that have completed their rumen development.



# SHELLMIN D3

**2500 ml**

Besides the product, it is supplied to the market with 25 liters of mineral mixture.



## COMPOSITION:

Each 1000 ml contains

Vitamin D3	20,000,000	IU
Excipients to	1,000	ml

**25 lt**

Besides the product, it is supplied to the market with 2.5 liters of Vitamin D3.



## COMPOSITION:

Each 1000 ml contains

Iron Chloride	2,500	mg	Sodium Chloride	10,000	mg
Zinc Chloride	2,500	mg	Sodium Selenite	50	mg
Manganese Chloride	2,500	mg	Magnesium Chloride Hexahydrate	7,250	mg
Copper Chloride	1,000	mg	Calcium Chloride	57,500	mg
Phosphoric Acid	30,000	mg	Excipients to	1,000	ml
Cobalt Chloride	22	mg			

## IT'S USED FOR:



## DOSAGE:

1000 ml Mineral Mixture + 100 ml Vitamin D3 / 1 ton of water is used.



## INDICATIONS:

- Mineral substance and Vitamin D3 are recommended to be used to meet their needs.
- Vitamin D3 is necessary for the proper absorption and use of calcium and phosphorus, which are necessary for normal growth, bone development and egg shell formation.
- Minerals are compounds involved in the metabolic process of eggshell formation. These trace minerals work as co-factors of enzymes involved in crust matrix formation. Carbonic anhydrase, which is zinc dependent, stimulates calcium carbonate deposition for eggshell formation. The manganese-dependent polymerase enzyme forms the shell glycoprotein matrix or foundation.
- Supplementing the diet with highly bioavailable minerals such as mineral-amino acid complexes increases eggshell weight and eggshell thickness. Copper affects the synthesis of the shell membrane by the activity of the copper-containing enzyme lysyl oxidase.
- Minerals play an important role in maintaining the osmotic pressure and thus regulate the exchange of water and solutes in the animal body.
- Minerals serve as structural components of soft tissues.
- Minerals are necessary for the transmission of nerve impulses and muscle contraction.
- Minerals play a vital role in the acid-base balance of the body and thus regulate the pH of the blood and other body fluids.







IT'S USED FOR:



# ADEPHOS



## COMPOSITIONS

Each 1000 ml contains:

Phosphorous (phosphoric Acid 75%)	120,000	mg
Calcium (Calcium Chloride 33.9%)	50,000	mg
Sodium (Sodium Chloride 39.3 %)	15,000	mg
Magnesium (Magnesium Chloride 11.9%)	11,000	mg
Copper (Copper Chloride 36.5%)	1,500	mg
Zinc (Zinc Chloride 30%)	8,000	mg
Manganese (Manganese Chloride 26.37%)	10,000	mg
Excipients to	1,000	ml



## BENEFITS:

- Poultry require a variety of minerals in their diet to maintain their health and productivity.
- Calcium and phosphorus are essential macrominerals for laying hens, with a requirement of 3.5-4% calcium, 0.3-0.4% available phosphorus, and 0.2% sodium in their diets for egg production.
- Zinc, manganese, and copper are trace minerals that are needed for bird health and nutrient utilization, as well as for skeletal and eggshell integrity.



## INDICATIONS:

- It helps to increase growth and production.
- Treatment and prevention of diseases caused by mineral salt deficiency in animals.
- It is used in stress situations.
- Malnutrition and absorption therapy.
- Minerals serve as structural components of soft tissues.
- Minerals are necessary for the transmission of nerve impulses and muscle contraction.



## DOSAGE:

- Poultry:

1000 ml per 1000 liters of drinking water for 3-5 days.



1000 ml

# ADEMIN SOL



## COMPOSITIONS

Each 1000 ml contains:

Iron	27,800	mg
Zinc	7,000	mg
Phosphor	5,185	mg
Cobalt	400	mg
Copper	5,850	mg
Excipients to	1,000	ml



## INDICATIONS:

- It helps to increase growth and production.
- Treatment and prevention of diseases caused by mineral salt deficiency in animals.
- It is used in stress situations.
- Malnutrition and absorption therapy.
- Used it is used to increase the appetite of animals.

## IT'S USED FOR:



## DOSAGE:

- Poultry:  
1000 ml / 1000 liters of drinking water.
- Calves and Lambs:  
10 ml / 100 kg body weight.





# CARVA MENT



## COMPOSITIONS

Each 1000 ml contains:

Menthol	30,000	mg
Carvacrol	60,000	mg
Thymol	10,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- To improve immunity against outbreaks of viral and bacterial diseases.
- Improved feed intake and feed conversion.
- Enhanced digestion, expanded productive performance.
- Improving egg production, shell quality, fecundity and hatchability.



## DOSAGE:

Poultry:

1000 ml per 1000 liters of drinking water for 3-5 days.

IT'S USED FOR:







# OREGO-MIX - 5



## COMPOSITIONS

Each 1000 ml contains:

Oregano Oil	50,000	mg
Excipients to	1,000	ml



## BENEFITS:

- Origanum vulgare is a natural, less toxic, residue-free feed supplement for poultry compared to other synthetic ingredients. It contains important bioactive components such as thymol and carvacrol.
- O.vulgare has antimicrobial, antioxidant, antiviral, immunomodulatory and antiparasitic effects as a poultry feed supplement.
- Potential advantages of using thyme extracts in poultry diets include improved feed intake and feed conversion, improved digestion, expanded productive performance, reduced disease incidence and economic losses.



## INDICATIONS:

- It is used to improve immunity and increase the body's resistance to diseases.
- It is used as an auxiliary treatment in cases of viral, bacterial and parasitic infections.
- It is used as an alternative treatment to antibiotics, especially in organic production.
- Due to its aromatic scent, it acts as an appetizer and increases feed consumption after illness.
- It is used as an aid in various stress situations.



## DOSAGE:

- Poultry: 500 ml per 1000 liters of drinking water.
- Cattle and Horses: 10 ml per 100 kg bodyweight for 3 - 5 days.
- Sheep, Goats and Calves: 10 ml per 100 kg bodyweight for 3 - 5 days.

## IT'S USED FOR:





1000 ml

# OKASAK



## COMPOSITIONS

Each 1000 ml contains:

Garlic Oil	50,000	mg
Thyme Oil	10,000	mg
Eucalyptus Oil	4,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- Protects against infections.
- Reduces coccidia and cryptosporos.
- Prevents appetite problems.
- Improves the feed cycle.
- Shows antibacterial and antiviral properties.
- Increases the secretion of bile from the liver.
- Regulates bowel functions.
- Stimulates the immune system.
- It is a strong bactericide, virus and protozoa killer, especially Salmonella and E.Coli.



## DOSAGE:

- Poultry: 500 ml per 1000 liters of drinking water.
- Cattle and Horses: 10 ml per 100 kg bodyweight for 3 - 5 days.
- Sheep, Goats and Calves: 10 ml per 100 kg bodyweight for 3 - 5 days.

## IT'S USED FOR:





1000 ml

#### IT'S USED FOR:



# IMMUNE STAR



## COMPOSITIONS

Each 1000 ml contains:

Maya Extract	100,000	mg
Artichoke Extract	20,000	mg
Citrus Extract	20,000	mg
Sorbitol	200,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- An immune booster used in poultry to improve resistance to viral and bacterial diseases.
- Use before and after vaccinations for best resistance.
- Prevention of Intestinal Diseases caused by pathogenic bacteria or fungi.
- Improving egg production, shell quality, fecundity and hatchability.



## DOSAGE:

- Poultry: 1000 ml per 1000 liters of drinking water.
- Cattle and Horses: 10 ml per 100 kg bodyweight for 3 – 5 days.
- Sheep, Goats and Calves: 10 ml per 100 kg bodyweight for 3 – 5 days.







1000 ml

# IMMUNEMAX



## COMPOSITIONS

Each 1000 ml contains:

Lysine	30,000	mg
L-Threonine	51,000	mg
DL-Methionine	6,600	mg
L-Tryptophan	20,000	mg
Citric Acid	70,000	mg
Lactic Acid 80%	110,000	mg
Zinc Glycinate	75,000	mg
Copper Glycinate	25,000	mg
Maya Extract	30,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- An immune booster used in poultry to improve resistance to viral and bacterial diseases.
- Use before and after vaccinations for best resistance.
- Prevention of Intestinal Diseases caused by pathogenic bacteria or fungi.
- Improving egg production, shell quality, fecundity and hatchability.



## DOSAGE:

- Poultry: 1000 ml per 1000 liters of drinking water for 3 – 5 days.

IT'S USED FOR:





# LIVOMIX



## COMPOSITIONS

Each 1000 ml contains:

Sorbitol	200,000	mg
L-Carnitine	50,000	mg
Choline Chloride 75%	25,000	mg
Magnesium Sulphate	200,000	mg
Vitamin C	10,000	mg
Artichoke Extract	1,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- LIVOMIX is hepato protective (liver protector), lipotropic (compounds that catalyze fat breakdown and transport) and detoxifying (components that ensure removal of toxins from the organism) agent. It detoxifies the liver and kidney tracts.
- Shows a regenerative and protective effect after treatments with drugs such as antibiotics, antiparasitic, anesthetics.
- It facilitates the digestion and metabolism of saturated fats by increasing the secretion of bile.
- Ensures well-balanced energy metabolism through their components involved in many enzymatic and biological reactions.
- Increases the quality of the carcass and ensures the production of meat that is free from toxins. It decreases the fat content of meat and increases the protein ratio.
- In breeders with liver disorders (especially fatty liver), it prevents ruptures in the liver and oviduct during ovulation, which sometimes have the risk of death.
- Strengthens the muscle and bone structure and optimizes muscle activities.
- Decreases the mortality rate by preventing the disorders of muscle-bone system, fat metabolism and stress-related disorders in new transport of young animals.
- Regulates the animal's energy metabolism, strengthen the immune system through different mechanisms and suppress stress due to extracts of artichoke.

## IT'S USED FOR:



## DOSAGE:

**For oral administration via feed or drinking water.**

- Poultry: 1000 - 2000 ml per 1000 liters of drinking water for 3 - 5 days.
- Cattle and Cow: 5 ml / 100 kg B.W for 3 - 5 days.
- Sheep, Goats and Calves: 10 ml / 100 kg B.W for 3 - 5 days.
- Horse and Foal: 7.5 ml / 100 kg B.W for 3 - 5 days.





# • LIVOMIX FORT



## COMPOSITIONS

Each 1000 ml contains:

Sorbitol	100,000	mg
L-Carnitine	2,000	mg
Choline Chloride	37,500	mg
Betaine	12,000	mg
Vitamin B1	1,000	mg
Vitamin B2	1,000	mg
Niacin	3,000	mg
Calcium-D-Pantothenate	1,000	mg
Vitamin B6	2,000	mg
Biotin	50	mg
Vitamin B12	100	mg
Methionine	1,000	mg
Lysine	1,000	mg
Artichoke Extract	20,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- Help to balance fat depositions, support prevention of liver dysfunctions.
- It will help correct the disturbances in case of an increased supply of free fatty acids in the blood.
- Prevention of liver damage from mycotoxins & avoid perosis.
- Minimised liver damage due to pathogens like bacteria, virus, fungi & to overcome hypertrophy of liver.
- Increased liver secretions for better digestion and boost liver function.
- Faster recovery from stress & diseases of all types.

## IT'S USED FOR:



## DOSAGE:

- Poultry: 1000 - 2000 ml per 1000 liters of drinking water for 3-5 days.
- Cattle: 5 ml / 100 kg body weight for 3-5 days.
- Sheep, Goats and Calves: 10 ml / 100 kg body weight for 3-5 days.
- Horse: 7.5 ml / 100 kg body weight for 3 - 5 days.







# HEPAMIX



## COMPOSITIONS

Each 1000 ml contains:

Sorbitol	100,000	mg
L-Carnitine	25,00	mg
Betaine HCl	5,000	mg
Choline Chloride	75,000	mg
Calcium-D-Pantothenate	10,000	mg
Magnesium Sulphate	50,000	mg
DL-Methionine	4,000	mg
Vitamin C	500	mg
Excipients to	1,000	ml



## INDICATIONS:

- HEPAMIX Is hepato protective (liver protector), lipotropic (compounds that catalyze fat breakdown and transport) and detoxifying (components that ensure removal of toxins from the organism) agent. It detoxifies the liver and kidney tracts.
- Shows a regenerative and protective effect after treatments with drugs such as antibiotics, antiparasitic, anesthetics.
- It facilitates the digestion and metabolism of

saturated fats by increasing the secretion of bile.

- Ensures well-balanced energy metabolism through their components involved in many enzymatic and biological reactions.
- Increases the quality of the carcass and ensures the production of meat that is free from toxins. It decreases the fat content of meat and increases the protein ratio.
- In breeders with liver disorders (especially fatty liver), it prevents ruptures in the liver and oviduct during ovulation, which sometimes have the risk of death.
- Strengthens the muscle and bone structure and optimizes muscle activities.
- Decreases the mortality rate by preventing the disorders of muscle-bone system, fat metabolism and stress-related disorders in new transport of young animals.



## DOSAGE:

**For oral administration via feed or drinking water.**

- Poultry: 1000 ml per 1000 liters of drinking water for 3 - 5 days.
- Cattle and Horses: 10 ml per 100 kg bodyweight for 3 - 5 days.
- Sheep, Goats and Calves: 20 ml per 100 kg bodyweight for 3 - 5 days.
- Mixed with feed, the product should be used immediately.

## IT'S USED FOR:





# DIUREMIX



## COMPOSITIONS

Each 1000 ml contains:

Magnesium Sulphate	120,000	mg
L-Carnitine	30,000	mg
Vitamin B1	4,000	mg
Niacin	5,000	mg
Vitamin B6	4,000	mg
Vitamin K	5,000	mg
Vitamin C	20,000	mg
DL-Methionine	10,000	mg
Lysine	12,000	mg
Betaine	20,000	mg
Potassium Citrate	30,000	mg
Excipients to	1,000	ml



## BENEFITS:

- L-carnitine has indispensable functions in intermediary metabolism. It plays an obligate role in fatty acid metabolism by directing fatty acids into the mitochondrial oxidative pathway through the action of specialized acyltransferases. In poultry production, L-carnitine has a multi-functional purpose, which includes: growth promotion, strengthening the immune system, antioxidant effects and improving semen quality.
- Diuretics are used to remove inappropriate water volume in animals with oedema or volume overload, correct specific ion imbalances, and

reduce blood pressure and pulmonary capillary wedge pressure.

- Amino acids and vitamins are important for preventing oxidative stress, regulating the immune response and maintaining normal physiological, biochemical and homeostatic mechanisms.

## IT'S USED FOR:



## INDICATIONS:

- Diuretic stimulant and regulator of kidney and liver functions in poultry (gout - various poisonings).
- Used to open the bird's appetite and provide it with vitality and activity.
- It is used in cases of compensation for nutritional deficiency in cases of various diseases and stress.
- It is used in the prevention of sudden death in birds due to the support of the liver and kidney functions.



## DOSAGE:

- Poultry:

350 ml / 1000 liters of drinking water for 3 – 5 days.



1000 ml

# • BITKI POWER



## COMPOSITIONS

Each 1000 ml contains:

Eucalyptus Oil	20,000	mg
Thyme Oil	20,000	mg
Menthol	30,000	mg
Excipients to	1,000	ml

that appear after vaccinations.

- It helps in the treatment of congestion of the mucous membranes of the airways.
- It is used as a powerful disinfectant for the respiratory system and intestinal tract.
- As an appetizer and improves digestion.



## DOSAGE:

- Poultry: 1000 ml per 1000 liters of drinking water for 3 – 5 days.
- Cattle and Horses: 10 ml per 100 kg bodyweight for 3 - 5 days.
- Sheep, Goats and Calves: 10 ml per 100 kg bodyweight for 3 - 5 days.



## INDICATIONS:

- Expectorant for the respiratory system for poultry and expectorant and an expanded air bronchium.
- Elimination of respiratory symptoms associated with respiratory diseases

## IT'S USED FOR:







1000 ml

#### IT'S USED FOR:



# BREATHMIX



## COMPOSITIONS

Each 1000 ml contains:

Carvacrol	40,000	mg
Thymol Oil	25,000	mg
Eucalyptus	12,000	mg
Rosemary	20,000	mg
Menthol	20,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- Expectorant for the respiratory system for poultry and expectorant and an expanded air bronchium.
- Elimination of respiratory symptoms associated with respiratory diseases that appear after vaccinations.
- It helps in the treatment of congestion of the mucous membranes of the airways.
- It is used as a powerful disinfectant for the respiratory system and intestinal tract.
- As an appetizer and improves digestion.



## DOSAGE:

- Poultry: 1000 ml per 1000 liters of drinking water for 3 - 5 days.
- Cattle and Horses: 10 ml per 100 kg bodyweight for 3 - 5 days.
- Sheep, Goats and Calves: 10 ml per 100 kg bodyweight for 3 - 5 days.





# ELECTROSOL



## COMPOSITIONS

Each 1000 ml contains:

Calcium Chloride	97,107	mg
Magnesium Chloride	39,181	mg
Sodium Chloride	5,080	mg
Potassium Chloride	1,906	mg
Iron Chloride	4,357	mg
Cobalt Chloride Hexahydrate	404	mg
Zinc Chloride	8,336	mg
Copper Chloride Pentahydrate	2,684	mg
Sodium Selenate 45%	239	mg
Phosphoric Acid 85%	158,200	mg
DL-Methionine	1,000	mg
L-Lysine Hcl	1,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- solution is used for Perosis, Cannibalism, Cage fatigue, stress, Alopecia, dehydration, bone weakness, metabolic defects that are results of the mineral substance deficiency.
- It can be used to infectious diseases, treating the metabolic disorders generating from mineral deficiencies.
- It increases development and egg shell quality in poultry.
- Copper in solution helps iron to be absorbed from the intestine.
- Copper deficiency caused anemia..

## IT'S USED FOR:



## DOSAGE:

- Poultry:

300 ml / 1000 liters of drinking water for 3 - 5 days.

- Calves and Lambs:

5 ml / 100 kg body weight 3 - 5 days.





**1000 ml**

#### IT'S USED FOR:



#### DOSAGE:

- 300-500 ml/day from 3 days before birth to 4 days after birth.
- Used orally.



#### INDICATIONS:

- It is used to provide glycogen substance.
- Negative energy balance, reproductive and liver problems for dairy cows before calving and after calving.
- The risk of ketosis in dairy cows.
- Used as a supplemental source for essential amino acids such as A-D3-E and fat-soluble vitamins.

# ANTIKET



#### COMPOSITIONS

Each 1000 ml contains:

Propylene Glycol	300,000	mg
Glycerine	665,000	mg
Vitamin A	300,000	IU
Vitamin D3	75,000	IU
Vitamin E	300	mg
Niacin	300	mg
Carnitine	5,000	mg
L-Lysine Hcl	43.25	mg
DL-Methionine	9.45	mg
L-Threonine	12.25	mg
Monosodium Glutamate	50	mg
L-Proline	31	mg
L-Glycine	25.75	mg
L-Leucine	24.25	mg
L-Valin	23	mg
L-Arginine	20.25	mg
Aspartic Acid	17.5	mg
L-Alanine	16.25	mg
L-Phenylalanine	15	mg
L-Isoleucine	13.5	mg
L-Tyrosine	8.1	mg
Excipients to	1,000	ml







1000 ml

# • ADETOX - S



## COMPOSITIONS

Each 1000 ml contains:

Phosphoric Acid 85%	68,000	mg
Citric Acid	95,000	mg
Lactic Acid 80%	60,000	mg
Malic Acid	25,000	mg
Tartaric Acid	12,000	mg
Vitamin B6	2,100	mg
Propylene Glycol	60,000	mg
Maya Extract	45,000	mg
Formic Acid 85%	10,000	mg
Propionic Acid	30,000	mg
Acetic Acid 80%	22,000	mg
Betaine	12,000	mg
Copper Sulphate	12,000	mg
L-Lysine Hcl	18,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- It is used as an anti-mycotoxin.
- Protecting the immune system and improving the immune response to various vaccines.
- It lowers the PH level and thus eliminates pathogenic bacteria and fungi in the gastrointestinal tract.
- It is used as a growth and performance booster and improves the fodder conversion rate.
- It helps reduce the use of antibiotics and does not interfere with any type of medicine.



## DOSAGE:

- Poultry: 1000 ml per 1000 liters of water.
- Cows and Sheep: 10 ml / 100 kg live weight per day.

## IT'S USED FOR:





IT'S USED FOR:



# ACID MIX



## COMPOSITIONS

Each 1000 ml contains:

Lactic Acid	40,000	mg
Phosphoric Acid	150,000	mg
Formic Acid	150,000	mg
Citric Acid	4,000	mg
Propionic Acid	10,000	mg
Acetic Acid	10,000	mg
Copper Glycinate	8,000	mg
Excipients to	1,000	ml



## INDICATIONS:


- Reduces PH of the gout
- Prevents colonization of pathogens like salmonella, E Coli & Clostridium.
- Maintains proper micro flora in the gout.
- Provides clean environment and healthy production
- Inhibits the growth of pathogenic intestinal microflora.
- This inhibition reduces the microflora competing for the host nutrients and results in better growth and performance of the chicken.



## DOSAGE:

- Poultry: 50 ml per 1000 liters of drinking water for 3-5 days.





# ANTI MOLD



## COMPOSITIONS

Each 1000 ml contains:

Citric Acid	25,000	mg
Lactic Acid	11,000	mg
Sorbic Acid	3,000	mg
Malic Acid	21,000	mg
Acetic Acid	70,000	mg
Propionic Acid	160,000	mg
Phosphoric Acid	22,000	mg
Formic Acid	75,000	mg
Excipients to	1,000	ml



## INDICATIONS:

- It is used as an anti-fungal
- It lowers the pH level and thus eliminates pathogenic bacteria and fungi in the gastrointestinal tract
- It is used as a growth and performance booster and improves the fodder conversion rate.
- It helps reduce the use of antibiotics and does not interfere with any type of medicine.



## DOSAGE:

- Poultry:  
1000 ml per 1000 liters of drinking water  
for 3 - 5 days.

## IT'S USED FOR:







# WATER SOLUBLE POWDER

ACK - MIX

BCK - MIX

C MIX 65%

C MIX 20%

E.S MIX W.S

ADE - MULTIVIT W.S.P

VITAMINO POWER

SUPER VIT PLUS

ANTI-STRESS FORTE

CALF-MIX SC

PROPIOMIX

SILAJ MIX

DIARESTAT

UREMIX

UREMIX FORT



# ACK - MIX



## COMPOSITIONS

Each 1000 gr contains:

Vitamin A	30,000,000	IU
Vitamin C	10,000	mg
Vitamin K3	25,000	mg
Excipients to	1,000	gr



## BENEFITS:

ACK - MIX provides water-soluble vitamins essential for carbohydrate and fat metabolism as well as protein synthesis.

ACK - MIX:

- Protecting nerve function, skin, hair and muscle tissue.
- Vital for all energy supply systems.
- Necessary for red blood cell formation.
- Helps reduce the effects of stress.
- Beneficial for foot ailments, paralysis and lameness.
- To increase the quality of the fleece and feather.



## INDICATIONS:

- Prevention or treatment of vitamin deficiencies in farm animals.
- Prevention or treatment of stress (caused by vaccination, illness, transportation, high humidity, high temperature or extreme temperature changes).
- Improvement of feed transitions.



## DOSAGE:

- Poultry: 1000 gr per 1000 liters of drinking water.
- Cattle and Horses: 10 gr per 100 kg bodyweight for 3 - 5 days.
- Sheep, Goats and Calves: 10 gr per 100 kg bodyweight for 3 - 5 days.



1000 gr

## IT'S USED FOR:







1000 gr

### IT'S USED FOR:



### DOSAGE:

- Poultry: 250 - 500 gr per 1000 liters of drinking water.
- Cattle and Horses: 10 gr per 100 kg bodyweight for 3 - 5 days.
- Sheep, Goats and Calves: 10 gr per 100 kg bodyweight for 3 - 5 days.

## BCK - MIX



### COMPOSITIONS

Each 1000 gr contains:

Vitamin B1	1,750	mg
Vitamin B2	3,500	mg
Niacin	17,500	mg
Calcium-D-Pantothenate	5,000	mg
Vitamin B6	1,750	mg
D-Biotin	35	mg
Vitamin B12	10	mg
Vitamin C	45,000	mg
Vitamin K3	3,000	mg
Excipient to	1,000	gr

- Necessary for red blood cell formation.
- Helps reduce the effects of stress.
- Beneficial for foot ailments, paralysis and lameness.
- Improving the quality of fleece.



### INDICATIONS:

- Prevention or treatment of vitamin B deficiencies in farm animals.
- Prevention or treatment of stress (caused by vaccination, illness, transportation, high humidity, high temperature or Zextreme temperature changes).
- Used to improve feed passage.



### BENEFITS:

BCK - MIX provides water-soluble vitamins essential for protein synthesis as well as carbohydrate and fat metabolism.

BCK - MIX:

- Protect nerve function, skin, hair and muscle tissue.
- Vital for all energy supply systems.







1000 gr

#### IT'S USED FOR:



## C MIX 65%



#### COMPOSITIONS

Each 1000 gr contains:

Vitamin C	650,000	mg
Excipients to	1,000	gr



#### BENEFITS:

• Ascorbic acid (Vitamin C) is a water-soluble antioxidant compound, which protects cells

against oxidative damage and improves immune system function.

• Vitamin C is believed to play a role in protein, lipid and carnitine synthesis, maintaining blood vessel integrity, and immune function.

• Vitamin C may also act as a co-antioxidant with other antioxidants by synergistic effects.

• Adding supplemental vitamin C to the diet is beneficial for birds subjected to stress, including heat stress from exposure to hot weather, Vitamin C will help enhance their immune system and alleviate metabolic signs of stress.



#### INDICATIONS:

• It used in poultry and ruminants in cases of heat stress induced problems such as poor immunity,

• Low feed intake, Low weight gain, oxidative stress, poor fertility and semen quality, low carcass weight and increase of mortality.



#### DOSAGE:

- Poultry: 250 gr per 1000 liters of drinking water for 5 days.

- Sheep and Goats: 1 gr each one in drinking water for 7 days.

- Calves: 10 gr each one in drinking water for 7 days.





1000 gr

#### IT'S USED FOR:



## C MIX 20%



#### COMPOSITIONS

Each 1000 gr contains:

Vitamin C	200,000	mg
Excipients to	1,000	gr



#### BENEFITS:

- Ascorbic acid (Vitamin C) is a water-soluble antioxidant compound, which protects cells against oxidative damage and improves immune system function.
- Vitamin C is believed to play a role in protein, lipid and carnitine synthesis, maintaining blood vessel integrity, and immune function.
- Vitamin C may also act as a co-antioxidant with other antioxidants by synergistic effects.
- Adding supplemental vitamin C to the diet is beneficial for birds subjected to stress, including heat stress from exposure to hot weather, Vitamin C will help enhance their immune system and alleviate metabolic signs of stress.



#### INDICATIONS:

- It used in poultry and ruminants in cases of heat stress induced problems such as poor immunity,
- Low feed intake, Low weight gain, oxidative stress, poor fertility and semen quality, low carcass weight and increase of mortality.



#### DOSAGE:

- Poultry: 1000 gr per 1000 liters of drinking water for 5 days.
- Sheep and Goats: 3 gr each one in drinking water for 7 days.
- Calves: 25 gr each one in drinking water for 7 days.





# E.S Mix W.S



1000 gr

## IT'S USED FOR:



## COMPOSITIONS

Each 100 gr contains:

Vitamin E	2,000	mg
Selenium	45	mg
Excipients to	100	gr



## SYMPTOMS OF A VITAMIN E DEFICIENCY INCLUDE:

- Crazy chick disease (encephalomalacia), Wry neck, Decreased laying Muscular dysfunction, Weakness, Inability to walk, bowed legs, Greenish -blue skin, Muscle spasms, Inability to control head (to turn it, hold it up or lower it) Hemorrhages, Paralysis, Enlarged hocks, Poor growth, Poor feed conversion Increase susceptibility to disease and parasites and Decreased fertility.



## BENEFITS:

- Vitamin E is a powerful antioxidant and important for normal neurological functions, Greater resistance to heat stress, strengthened immune response, including to E. coli, Improved laying, A better hatch rates, Higher levels of vitamin E in eggs and meat.
- Selenium has been shown to have a strong antioxidant, anti-inflammatory, and antimicrobial ability.



## INDICATIONS:

- It is used in cases of vitamin E deficiency in calves, lambs, sheep, goats, piglets and poultry.
- Encephalomalacia (Crazy Chick Disease).
- Muscular dystrophy (White Muscle Disease, Stiff Lamb Disease).
- Decreased hatchability of eggs.
- Dietary selenium supplementation has been used for improving growth performance, immune function, and meat quality.
- Selenium is an essential trace mineral important for the maintenance of homeostasis in animals.



## DOSAGE:

- Poultry: 1000 gr per 1000 liters of drinking water 3-5 days.
- Livestock: 10 gr per 100 kg body Wight for 3-5 days.





# ADE - MULTIVIT W.S.P



1000 gr

## IT'S USED FOR:



## INDICATIONS:

- ADE MULTIVIT is used as a vitamin and amino acid supplement.
- Fat - soluble vitamins such as A - D3 - E.
- Water - soluble vitamins such as B vitamin complexes and vitamin C.
- It is used in cases of vitamin deficiency, keeping the birds healthy, increasing production, increasing fertility and helping the formation of immunity.



## COMPOSITIONS

Each 1000 gr contains:

Vitamin A	10,000,000	IU
Vitamin D3	2,000,000	IU
Vitamin E	2,000	mg
Vitamin B1	50	mg
Vitamin B2	4	mg
Niacin	1,000	mg
Calcium-D-Pantothenate	100	mg
Vitamin B6	500	mg
Biotin	10,000	mcg
Folic Acid	10	mg
Vitamin B12	1,000	mcg
Vitamin C	2,000	mg
Vitamin K3	10	mg
Choline Chloride	350	mg
Magnesium	5,000	mg
Zinc	5,000	mg
Iron	1,000	mg
Cobalt	100	mg
Selenium	110	mg
DL-Methionine	1,000	mg
L-Lysine Hcl	1,000	mg
L-Threonine	500	mg
Betaine	100	mg
Excipients to	1,000	gr



## DOSAGE:

- Poultry: 1000 gr per 1000 liters of drinking water.
- Cattle and Horses: 10 gr per 100 kg bodyweight for 3 - 5 days.
- Sheep, Goats and Calves: 10 gr per 100 kg bodyweight for 3 - 5 days.



# VITAMINO POWER



1000 gr

## IT'S USED FOR:



## COMPOSITIONS

Each 1000 gr contains:

Vitamin A	8,000,000	IU
Vitamin D3	1,000,000	IU
Vitamin E	10,000	mg
Vitamin B1	1,000	mg
Vitamin B2	1,000	mg
Niacin	15,000	mg
Calcium-D-Pantothenate	4,000	mg
Vitamin B6	1,000	mg
Biotin	15,000	mcg
Folic Acid	200	mg
Vitamin B12	5,000	mcg
Vitamin C	7,000	mg
L - Threonine	2,450	mg
L - Valin	4,600	mg
L - Alanine	3,250	mg
L - Arginine	4,050	mg
Aspartic Acid	3,500	mg
L - Phenylalanine	3,000	mg
L - Glycine	5,150	mg
Monosodium Glutamate	7,050	mg
L - Isoleucine	2,700	mg
L - Leucine	4,850	mg
L - Lysine Hcl	8,650	mg
DL-Methionine	1,890	mg
L - Proline	6,200	mg
L - Tyrosine	1,620	mg
Iron	8,000	mg
Zinc	7,250	mg
Manganese	9,300	mg
Copper	1,250	mg
Selenium	100	mg
Excipients to	1,000	gr



## INDICATIONS:

- VETAMINO POWER is used as a supplementary source of poultry, essential amino acids, fat-soluble vitamins such as A-D3-E, and water-soluble vitamins such as B-vitamin complexes and vitamin C.
- It also contains amino acids and minerals.



## DOSAGE:

- Poultry: 1000 gr per 1000 liters of drinking water for 3 - 5 days.
- Calves and lambs: 10 g per 100 kg body weight daily.
- Horses: 10 g per 100 kg body weight 3-5 days before the race.





# SUPER VIT PLUS



1000 gr

## IT'S USED FOR:



## COMPOSITIONS

Each 1000 gr contains:

Vitamin A	8,000,000	IU
Vitamin D3	1,500,000	IU
Vitamin E	6,000	mg
Vitamin B1	1,250	mg
Vitamin B2	2,000	mg
Niacin	12,500	mg
Vitamin B6	1,000	mg
Folic Acid	400	mg
Vitamin B12	5,000	mcg
Vitamin K3	2,000	mg
Vitamin C	1,500	mg
L - Threonine	1,225	mg
L - Valin	2,300	mg
L - Alanine	1,625	mg
L - Arginine	2,025	mg
L - Aspartic Acid	1,750	mg
L - Phenylalanine	1,500	mg
L - Glycine	2,575	mg
Monosodium Glutamate	3,525	mg
L - Isoleucine	1,350	mg
L - Leucine	2,425	mg
L - Lysine Hcl	4,325	mg
DL-Methionine	945	mg
L - Proline	3,100	mg
L - Tyrosine	810	mg
Iron Sulphate	25,000	mg
Cobalt Sulphate	150	mg
Manganese Sulphate	15,000	mg
Zinc Sulphate	10,000	mg
Calcium Chloride	10,000	mg
Excipients to	1,000	gr



## INDICATIONS:

• Super vit plus is used as a supplementary source of essential amino acids, fat-soluble vitamins such as A-D3-E, and water-soluble vitamins such as B-vitamin complexes and vitamin C, in poultry, horses, calves and lambs.



## DOSAGE:

- Poultry:  
1000 gr per 1000 liters of drinking water for 3 - 5 days.

- Calves and Lambs:  
10 g per 100 kg body weight daily.

- Horses:  
10 g per 100 kg body weight 3 - 5 days before the race.





1000 gr

### IT'S USED FOR:



# ANTI-STRESS FORTE



## COMPOSITIONS

Each 1000 gr contains:

Salicylic Acid	250,000	mg
Caffeine	33,000	mg
Vitamin C	60,000	mg
Magnesium Sulphate	7,100	mg
Sodium Bicarbonate	50,000	mg
Sodium Chloride	70,000	mg
Sodium Sulphate	225,000	mg
Potassium Chloride	11,000	mg
Excipients to	1,000	gr



## BENEFITS:

Anti-stress forte Composition, Salicylic Acid, Vitamin C, Caffeine.

It is a feed additive in powder form containing a balanced combination of Sodium Chloride, Sodium Bicarbonate, Magnesium Sulphate, Potassium Chloride and Sodium Sulphate.



## INDICATIONS:

- It is used to effectively return the body to its optimum condition, especially in heat-related stress situations, environmental factors where body temperature increases, and infection situations.
- It eliminates the clinical effects of heat stress, lowers the rising body temperature, and helps animals to regain their optimum metabolism by reducing the response to stress.



## DOSAGE:

- Poultry: 500 gr / 1000 liters of drinking water for 3 - 5 days.
- Cows and Sheep: 10 gr / 100 kg body weight daily.
- Horses: 10 gr / 100 kg body weight 3 - 5 days before the race.

# CALF-MIX SC



## COMPOSITIONS

Each 10 kg contains:

Vitamin A	20,000,000	IU
Vitamin D3	3,000,000	IU
Vitamin E	25,000	mg
Vitamin B1	8,000	mg
Vitamin B2	4,000	mg
Niacin	20,000	mg
Calcium-D-Pantothenate	12,000	mg
Vitamin B6	5,000	mg
Biotin	50	mg
Vitamin B12	20	mg
Manganese	50,000	mg
Iron	50,000	mg
Zinc	50,000	mg
Copper	10,000	mg
Copper Glycinate	5,000	mg
Manganese Glycinate	10,000	mg
Zinc Glycinate	20,000	mg
Iodine	800	mg
Cobalt	150	mg
Selenium	150	mg
Saccharomyces Cerevisiae	100 × 10 <sup>11</sup>	CFU
Cholin Chloride 60%	96,000	mg
Pectin	100,000	mg
Kaolin	150,000	mg
Excipients to	10,000	gr



## DOSAGE:

- Small calves: It is given daily by mixing 10-15 Gr. Of milk and feed per calf.
- Or as veterinarian recommendations.



## INDICATIONS:

### For dairy and fattening calves

- It is a mixture containing high-density vitamin supplement. Biotin. Niacin. Live yeast and organic minerals in order to meet the vitamin and trace mineral substance needs of calves.
- The most important factor determining the profitability of cattle enterprises is to raise healthy calves.
- CALF MIX SC is a formula enriched with live yeast along with vitamins. Organic and inorganic minerals that calves need.
- Increases the calf's Growth rate. And is effective in solving respiratory tract problems.
- It supports beneficial bacteria in the digestive system and increases feed efficiency.
- Reduces diarrhea and prevents toxicity.
- CALF - MIX SC which helps you prevent calf deaths. Also minimizes the impact of calves during stress periods.



5 kg

## IT'S USED FOR:



# PROBIOMIX



## COMPOSITIONS

Each 1000 gr contains:

Lactobacillus Casei	60 x 10 <sup>8</sup>	CFU
Lactobacillus Acidophilus	240 x 10 <sup>8</sup>	CFU
Lactobacillus Plantarum	272 x 10 <sup>8</sup>	CFU
Lactobacillus Paracasei	392 x 10 <sup>8</sup>	CFU
Excipients to	1,000	gr



## WHAT IT DOES, HOW TO USE IT:

• Probiotics are beneficial bacteria and yeasts that help various body systems, especially the digestive system and skin, to work in a balanced way,

unlike bacteria that cause infections and diseases.

• Probiotics, especially probiotics in the gut, are largely aimed at ensuring the proper functioning of the digestive and immune systems. Beneficial bacteria play a major role in maintaining the intestinal flora balance.

### Endikasyonları: FOR ALL POULTRY

1. Contributes to the support of the immune system.
2. It helps digest food.
3. It helps the production of vitamins synthesized in the body.
4. It helps the cells that make up the intestinal wall and prevent harmful bacteria entering.
5. It increases the absorption of drugs taken from the outside into the body.
6. Helps maintain the physiological balance in the intestine.
7. Eliminates bloating problems.
8. Protects liver and bone health.
9. Controls endemic and zoonotic agents by competitive exclusion.
10. It slows down the infectious agents that cause deadly diseases (such as E.coli, Salmonella, Campylobacter).
11. It provides an increase in egg and fertility.



## DOSAGE:

- Unless recommended otherwise by the veterinarian.
- 500 gr is used for 1 ton of drinking water.



500 gr

## IT'S USED FOR:







500 gr

## IT'S USED FOR:



## APPLICATION RATES:

- One package (250 GR). It is used for 100 tons of corn. One package is diluted in 200 - 300 liters of water and used in 100 tons of silage. 1 - 2L/ton is sprayed.
- It can be used in all spraying systems. In Package Silages, 500.000 CFU buffer is applied to 1 gr. green plant in accordance with dry matter.

# SILAJ MIX



## COMPOSITIONS

Each 1000 gr contains:

Lactobacillus Plantarum	280 x 10 <sup>8</sup>	CFU
Lactobacillus Casei	66 x 10 <sup>8</sup>	CFU
Lactobacillus Acidophilus	280 x 10 <sup>8</sup>	CFU
Lactobacillus Paracasei	400 x 10 <sup>8</sup>	CFU
Xylanase Endo-1.4 Beta Xylanase	1,950,000	IU
Beta Glucanase	303,800	IU
Amylase Alpha Amylase	173,600	IU
Protease	303,800	IU
Excipients to	1,000	gr



## INDICATIONS:

- Provides Fast Fermentation. It reduces dry matter loss.
- Decreases protein breakdown. It increases its digestibility.
- Controls the heating and losses on the surface during silage intake and minimizes it.
- Produces antifungal propionic acid (kills yeast and mold).
- It enables us to obtain high quality silage rich in nutritional value.
- It helps the silage to be stored for a long time after opening without losing its quality.
- Thanks to the enzymes it contains, it accelerates the fermentation of cellulose and polysaccharides in the plant cell wall and contributes to the formation of lactic acid.



## HOW DOES IT WORK?

- During the silage making phase, the pH level of the mown plant has to drop from 6.5 to 4 as quickly as possible. It is to accelerate the fermentation of cellulose and similar polysaccharides in plant cell walls, to contribute to the formation of lactic acid, and to increase the digestibility of silage at the same time.
- The SILAGE MIX ingredient is a combination of selected elements to achieve a fast and effective fermentation.

# DIARESTAT



## COMPOSITIONS

Each 1000 gr contains:

Vitamin A	2,500,000	IU
Vitamin K3	9,000	mg
Vitamin C	10,000	mg
Copper	1,500	mg
Selenium	100	mg
Pectin	60,677	mg
Kaolin	900,000	mg
Excipients to	1,000	gr



## INDICATIONS:

- It is effective in non-specific diarrhea treatment.
- It plays a supportive role in the treatment of poisoning and its cases due to its astringent and surface-retaining activity in the digestive system.

- Due to the ACK vitamins it

contains, it also provides rapid healing of the destroyed intestines.

- It is used for the purpose of benefiting from feed efficiency and providing live weight gain.



## DOSAGE:

- Cow and Camel: 10 gr / Day.
- Calf and Foal: 5 gr / Day.
- Sheep and Goat: 5 gr / Day.
- Lamb and Capricorn: 3 gr / Day.

## DIARESTAT

### BİLEŞENLER:

Her 1000 gr için:

Vitamin A	2.500.000	IU
Vitamin K3	9.000	mg
Vitamin C	10.000	mg
Bakır	1.500	mg
Selenyum	100	mg
Pektin	60.677	mg
Kaolin	900.000	mg

### KULLANIM

- Herhangi bir türü sağlıklarında zayıflık
- Sindirim sistemlerinde bozuklukları ve sindirim
- Kuluçca etkenleri nedeniyle pat. mikroflora ve
- etkiyle oluşan enfeksiyonların önlenmesi için

- İçerdiği ACK vitaminlerinden dolayı aynı zamanda
- sindirim sistemi için destekleyici etkiyi de sağlar

- Özellikle zayıflıkta etkili olduğu ve sindirim
- sisteminde etkiyi artırarak kullanılır.

### DOZAJ:

- Tıbbi tedavilerde kullanılır: 10 gr / gün
- Sığır - Keçi - 10 gr / gün
- Koyun - 5 gr / gün
- Kuzu - 3 gr / gün
- Kedi - 0,5-1 gr / gün

Adetleri için kullanılır: 10 gr / gün

Her 1000 gr için:



100GR

### UYARILAR:

Her 1000 gr için 24 saat süreyle

kullanılır.

### DEPOLAMA ŞEKLİ:

Her 1000 gr için 24 saat süreyle

kullanılır.

Her 1000 gr için 24 saat süreyle

kullanılır.

100 gr

## IT'S USED FOR:



# UREMIX



## COMPOSITIONS

Each 1000 gr contains:

Ammonium Chloride	300,000	mg
Sodium Citrate	5,000	mg
Potassium Chloride	20,000	mg
Magnesium Sulphate	100,000	mg
Vitamin C	100,000	mg
DL-Methionine	100,000	mg
Betaine	100,000	mg
Excipients to	1,000	gr



## INDICATIONS:

- UREMIX is very effective in case of hydronephrosis and nephritis.
- UREMIX is also effective in case of infectious bursal disease, stimulant of hepatic functions and as a diuretic.
- UREMIX contains Ammonium chloride as a urinary acidifying agent to prevent and dissolve certain type of uroliths and to enhance renal excretion of some toxins.
- Potassium chloride and magnesium sulfate to mitigate the harmful effect of heat stress on birds to improve their productive performance.
- Methionine is an essential amino acid for growth and also as a urine acidifier.
- Vitamin C is essential for growth and for immunity enhancement.



## DOSAGE:

- Poultry:

1000 gr / 1000 liters of drinking water for  
2 - 3 days or 2 kg/ton of feed for 2-3 days.



1000 gr

## IT'S USED FOR:





# UREMIX FORT



## COMPOSITIONS

Each 1000 gr contains:

Ammonium Chloride	650,000	mg
DL-Methionine	100,000	mg
Betaine	50,000	mg
Vitamin C	75,000	mg
Vitamin A	500,000	IU
Excipients to	1,000	gr



## INDICATIONS:

- UREMIX FORT is very effective in case of hydronephrosis and nephritis.
- UREMIX FORT is also effective in case of infectious bursal disease, stimulant of hepatic functions and as a diuretic.
- UREMIX FORT contains Ammonium chloride as a urinary acidifying agent to prevent and dissolve certain type of uroliths and to enhance renal excretion of some toxins.
- Methionine is an essential amino acid for growth and also as a urine acidifier.
- Vitamins A and C are essential for growth and for immunity enhancement.



## DOSAGE:

- Poultry:

1000 gr / 1000 liters of drinking water for 2 – 3 days  
or 2 kg/ton of feed for 2-3 days.



1000 gr

## IT'S USED FOR:





# ANTITOXIN + ANTIOXIDANT PREMIXS

TOXSTAT PLUS

TOXSTAT

MYCOLINK


MYCOSTOP

TOXSTOP

SALKIL

SUPER OXISTOP

OXISTOP



# TOXSTAT PLUS



## COMPOSITIONS

Each 1 kg contains:

Organic Acids, Calcium Propionate, Calcium Formate,  
Mannan Oligosaccharide, Beta glucan, Hydrated Sodium  
Potassium Calcium Aluminium Silicate (HSCAS)



## INDICATIONS:

- Improves animal fertility and performance.
- Supports animal immune and antioxidant systems.
- Effective protection against a broad-spectrum of mycotoxins.



## DOSAGE:

0.5 – 1 Kg for 1 Ton of Feed.



25 kg



IT'S  
USED FOR  
CHICKENS







25 kg

 IT'S  
USED FOR  
CHICKENS

# TOXSTAT



## COMPOSITIONS

Each 1 kg contains:

Calcium Format	85,000	mg
Inactive Yeast	15,000	mg
Citric Acid	5,000	mg
Sorbic Acid	2,000	mg
Propionic Acid	5,000	mg
Sodium Bentonite	250,000	mg
Clinoptilolite	638,000	mg




## INDICATIONS:

- Provides effective protection against a broad spectrum of mycotoxins.
- Supports the immune and antioxidant systems of animals.
- It improves the fertility and performance of animals.



## DOSAGE:

- 1 kg per ton of feed.



# MYCOLINK



## COMPOSITIONS

Each 1 kg contains:

Inactive Yeast	10,000	mg
Sodium Bentonite	100,000	mg
Clinoptilolite	890,000	mg



## INDICATIONS:

- Provides effective protection against a broad spectrum of mycotoxins.
- Supports the immune and antioxidant systems of animals.
- It improves the fertility and performance of animals.



## DOSAGE:

1 kg per ton of feed.



IT'S  
USED FOR  
CHICKENS







25 kg

# MYCOSTOP



## COMPOSITIONS

Each 1 kg contains:

Zeolite, Milk Thistle, Bentonite, Mannan Oligosaccharide  
Beta Glucan, Hydrated Sodium Calcium Aluminium Silicate (HSCAS)



## INDICATIONS:

- Fights a wide range of mycotoxins (enzymatic deactivation of trichothecenes, zearalenone, ochratoxin A and fumonisins).
- It supports the immune and antioxidant systems of animals.
- It improves the fertility and performance of animals.
- Highest aflatoxin adsorption.
- Advanced biological protection.
- Protection against endotoxins.



## DOSAGE:

- 0.5 – 1 Kg for 1 Ton of Feed.

## IT'S USED FOR:







25 kg

#### IT'S USED FOR:



## TOXSTOP



#### COMPOSITIONS

Each 1 kg contains:

Inulin	30,000	mg
Active Molecule Silymarin	60,000	mg
Thyme Oil	2,000	mg
Active Carbon	30,000	mg
Clinoptilolite	600,000	mg
Malt Grass	278,000	mg



#### INDICATIONS:


- Provides effective protection against a broad spectrum of mycotoxins.
- Causes changes in the gut microbiota by selectively promoting healthy positive bacterial strains.
- Supports the immune and antioxidant systems of animals.
- Increasing the secretion of digestive enzymes, reducing the viscosity of digestive materials, strengthening the immune system, lowering cholesterol in the blood.
- It has antioxidant and antiviral properties.
- It helps prevent the growth of fungi, purifies animals from diseases and stimulates the appetite of animals.



#### DOSAGE:

(0.5 - 2) kg per ton of feed.





25 kg

IT'S  
USED FOR  
CHICKENS

# SALKIL



## COMPOSITIONS

Each 1 kg contains:

Ammonium Format	250,000	mg
Ammonium Propionate	74,000	mg
Formic Acid 85%	80,000	mg
Clinoptilolite	556,000	mg



## INDICATIONS:

- Contains formic acid (1K236) and other organic acids with known antimicrobial activity to help control Campylobacter, Salmonella, Escherichia coli and other Gram negative microorganisms in raw materials and processed feeds.
- Maintains a balanced and diverse gut microbiota.
- Supports normal digestive processes and the production of internal enzymes necessary to maintain healthy growth and performance.
- Helps feed acidification, increases flavor and digestibility.
- Supports a strong stomach barrier necessary to maintain general health.
- It helps to prevent mold and bacteria formation in feed.



## DOSAGE:

1 kg per ton of feed.





# SUPER OXISTOP



25 kg

 IT'S  
USED FOR  
CHICKENS



## COMPOSITIONS

Each 1 kg contains:

Butylated Hydroxy Toluene (BHT)	200,000 mg
Butylated Hydroxy Anisole (BHA)	200,000 mg



## INDICATIONS:

- Antioxidant to be added to animals and fish premixes and feed.



## DOSAGE:

- 25 - 50 gr per 1 ton of final feed.







25 kg

IT'S  
USED FOR  
CHICKENS

# OXISTOP



## COMPOSITIONS

Each 1 kg contains:

Butylated Hydroxy Toluene (BHT)	110,000 mg
Butylated Hydroxy Anisole (BHA)	10,000 mg
Sodium Citrates	50,000 mg



## INDICATIONS:

- Antioxidant to be added to animals and fish premixes and feed.



## DOSAGE:

125 g per 1 ton of final feed.





# POULTRY PREMIXES

ADEMIX VITAMIN LAYER BRE 0.1%

ADEMIX VITAMIN LAYER PRO 0.1%

ADEMIX MINERAL LAYER 0.1%

ADEMIX LAYER 1%

ADEMIX LAYAR 2.5%

SUPER SHELL

VITAMIN BROILER STARTER 0.1%

VITAMIN BROILER GROWER 0.1%

VITAMIN BROILER FINISHER 0.1%


MINERAL BROILER 0.1%

ADEMIX BROILER STARTER 2.5%

ADEMIX BROILER GROWER 2.5%

ADEMIX BROILER FINISHER 2.5%

PROBIOFEED







# OUR PREMIXES PRODUCTS

---

## POULTRY


Why are feed additives needed to provide extra nutrition?

It has been known for a long time that general raw feed composed of basic raw materials such as soybeans, corn and fat does not contain all the necessary nutrients, which makes the feed insufficient for the animal to obtain the best performance. Another fact is that problems such as low digestibility, microbial contamination, and tarnished color can be fixed by adding ingredients such as vitamins, enzymes, and amino acids are added to livestock feed. These two facts above are just two normal examples where feed additives are useful and helpful. To speak in a sum way, feed additives are the key to help animals to get complete nutrition and therefore have a healthier body condition. Since the relationship between the health of humans and the health of animals is very close, helping animals to become the best of themselves is helpful for humans as well.

The main reason why feed additives are used so often in farm industries nowadays is that they are equipped with a great variety of nutrition. These special nutritions not only help to give animals the health they want but also help to maximize the nutrition of feedstuff for animals, thereby better the production efficiency in farm industries.







25 kg



## ADEMIX VITAMIN LAYER BRE 0.1%



### COMPOSITIONS

Each 1 kg contains:

Vitamin A	10,000,000 IU
Vitamin D3	2,000,000 IU
Vitamin E	25,000 mg
Vitamin B1	1,000 mg
Vitamin B2	6,000 mg
Niacin	30,000 mg
Calcium D-Pantothenate	8,000 mg
Vitamin B6	3,000 mg
Biotin	50,000 mcg
Folic Acid	1,000 mg
Vitamin B12	20,000 mcg
Vitamin K3	3,000 mg
Anti-Caking	Included
Anti-Oxidant	Included




### DOSAGE:

1 Kg for 1 Ton of Feed.



## ADEMIX VITAMIN LAYER PRO 0.1%



25 kg

 IT'S  
USED FOR  
CHICKENS



### COMPOSITIONS

Each 1 kg contains:

Vitamin A	10,000,000	IU
Vitamin D3	2,500,000	IU
Vitamin E	25,000	mg
Vitamin B1	1,000	mg
Vitamin B2	4,000	mg
Niacin	30,000	mg
Calcium D-Pantothenate	10,000	mg
Vitamin B6	3,000	mg
Biotin	50,000	mcg
Folic Acid	500	mg
Vitamin B12	25,000	mcg
Vitamin K3	3,000	mg
Anti-Oxidant	Included	
Anti-Caking	Included	




### DOSAGE:

1 Kg for 1 Ton of Feed.







25 kg

IT'S  
USED FOR  
CHICKENS

## ADEMIX MINERAL LAYER 0.1%



### COMPOSITIONS


Each 1 kg contains:

Manganese	100,000 mg
Copper	5,000 mg
Zinc	60,000 mg
Iron	25,000 mg
Iodine	500 mg
Selenium	200 mg




### DOSAGE:

1 Kg for 1 Ton of Feed.







 IT'S  
USED FOR  
CHICKENS



## ● ADEMIX LAYER 1%



### COMPOSITIONS

Each 10 kg contains:


Vitamin A	10,000,000	IU
Vitamin D3	3,000,000	IU
Vitamin E	25,000	mg
Vitamin B1	2,000	mg
Vitamin B2	5,000	mg
Niacin	40,000	mg
Calcium D-Pantothenate	12,000	mg
Vitamin B6	5,000	mg
Biotin	50	mg
Folic Acid	750	mg
Vitamin B12	20	mg
Vitamin K3	3,000	mg
Vitamin C	100,000	mg
Choline Chloride	100,000	mg
Phytase	400,000	FUT
Xylanase	1,275,000	FUT
Manganese	100,000	mg
Iron	60,000	mg
Zinc	60,000	mg
Copper	8,000	mg
Iodine	1,000	mg
Selenium	250	mg
Cobalt	150	mg
BHT	30,000	mg
BHA	7,500	mg
Phosphor	4.5	%
Calcium	16	%
Sodium	4	%
DL-Methionine	15	%
L-Lysine	7.5	%
Antioxidant	Included	
Anti-Caking	Included	



### DOSAGE:

10 Kg for 1 Ton of Feed.





25 kg



## ADEMIX LAYER 2.5%



### COMPOSITIONS

Each 1 kg contains:

Vitamin A	440,000 IU	L-Lysine Hcl	2 %
Vitamin D3	110,000 IU	Calcium	22 %
Vitamin E	1,400 mg	Phosphor	4.5 %
Vitamin B1	100 mg	Phytase	16,000 FUT
Vitamin B2	160 mg	Xylanase	51,000 FUT
Niacin	1,200 mg	Antitoxin	Included
Calcium D-Pantothenate	400 mg	Anti-Caking	Included
Vitamin B6	120 mg	Anti-Oxidant	Included
Biotin	2,000 mcg		
Folic Acid	21 mg		
Vitamin B12	1,000 mcg		
Vitamin K3	120 mg		
Choline Chloride 60%	30,000 mg		
Manganese	4,010 mg		
Iron	1,600 mg		
Zinc	2,420 mg		
Copper	600 mg		
Iodine	40 mg		
Selenium	10 mg		
Sodium Chloride	80,000 mg		
Bicarbonate Sodium	80,000 mg		
DL-Methionine	5.55 %		



### DOSAGE:

25 Kg for 1 Ton of Feed.







25 kg

IT'S  
USED FOR  
CHICKENS



## SUPER SHELL



### COMPOSITIONS

Each 1 kg contains:

Vitamin D3	2,500,000	IU
Copper (Copper Glycinate)	5,000	mg
Zinc (Zinc Glycinate)	30,000	mg
Manganese (Manganese Glycinate)	30,000	mg
Selenium (Selenium Glycinate)	50	mg



### INDICATIONS:

- Vitamin D3 is required for proper absorption and utilization of calcium and phosphorus, which are required for normal growth, bone development, and eggshell formation.
- Zinc, Manganese and copper are compounds involved in the metabolic process of eggshell formation. These trace minerals work as co-factors of enzymes involved with shell matrix formation. Carbonic anhydrase, which is zinc dependant, stimulates calcium carbonate deposition for eggshell formation. Polymerase enzyme, which is dependent on manganese, forms the shell glycoprotein matrix or foundation.
- Supplementing the diet with highly bioavailable minerals like mineral-amino acid complexes increases the eggshell weight and eggshell thickness. Copper affects the synthesis of shell membrane by activity of copper containing enzyme lysyl oxidase.
- Minerals play a key role in the maintenance of osmotic pressure, and thus regulate the exchange of water and solutes within the animal body.
- Minerals serve as structural constituents of soft tissues.
- Minerals are essential for the transmission of nerve impulses and muscle contraction.
- Minerals play a vital role in the acid-base equilibrium of the body, and thus regulate the pH of the blood and other body fluids.



### DOSAGE:

1 Kg for 1 Ton of Feed.





25 kg

 IT'S  
USED FOR  
CHICKENS

## ADEMIX VITAMIN BROILER STARTER 0.1%



### COMPOSITIONS

Each 1 kg contains:

Vitamin A	13,000,000 IU
Vitamin D3	5,000,000 IU
Vitamin E	80,000 mg
Vitamin B1	5,000 mg
Vitamin B2	9,000 mg
Niacin	70,000 mg
Calcium D-Pantothenate	25,000 mg
Vitamin B6	5,000 mg
Biotin	350,000 mcg
Folic Acid	2,500 mg
Vitamin B12	20,000 mcg
Vitamin K3	4,000 mg
Anti-Oxidant	Included
Anti-Caking	Included



### DOSAGE:

1 Kg for 1 Ton of Feed.





25 kg

IT'S  
USED FOR  
CHICKENS

## ADEMIX VITAMIN BROILER GROWER 0.1%



### COMPOSITIONS

Each 1 kg contains:


Vitamin A	11,000,000 IU
Vitamin D3	4,500,000 IU
Vitamin E	65,000 mg
Vitamin B1	4,000 mg
Vitamin B2	8,000 mg
Niacin	65,000 mg
Calcium D-Pantothenate	20,000 mg
Vitamin B6	4,000 mg
Biotin	280,000 mcg
Folic Acid	2,000 mg
Vitamin B12	18,000 mcg
Vitamin K3	3,600 mg
Anti-Oxidant	Included
Anti-Caking	Included



### DOSAGE:

1 Kg for 1 Ton of Feed.





25 kg

IT'S  
USED FOR  
CHICKENS

## ADEMIX VITAMIN BROILER FINISHER 0.1%



### COMPOSITIONS

Each 1 kg contains:

Vitamin A	10,000,000 IU
Vitamin D3	4,000,000 IU
Vitamin E	55,000 mg
Vitamin B1	3,000 mg
Vitamin B2	7,000 mg
Niacin	50,000 mg
Calcium D-Pantothenate	15,000 mg
Vitamin B6	3,000 mg
Biotin	220,000 mcg
Folic Acid	1,800 mg
Vitamin B12	16,000 mcg
Vitamin K3	3,200 mg
Anti-Oxidant	Included
Anti-Caking	Included




### DOSAGE:

1 Kg for 1 Ton of Feed.







25 kg

 **IT'S  
USED FOR  
CHICKENS**

## ADEMIX MINERAL BROILER 0.1%



### COMPOSITIONS

Each 1 kg contains:

Manganese	120,000	mg
Copper	16,000	mg
Zinc	120,000	mg
Iron	20,000	mg
Iodine	1,250	mg
Selenium	300	mg
Anti-Caking	Included	



### DOSAGE:

1 Kg for 1 Ton of Feed.





25 kg



## ADEMIX BROILER STARTER 2.5%



### COMPOSITIONS

Each 1 kg contains:


Vitamin A	520,000	IU
Vitamin D3	200,000	IU
Vitamin E	3,200	mg
Vitamin K3	160	mg
Vitamin B1	200	mg
Vitamin B2	360	mg
Niacin	2,800	mg
Calcium D-Pantothenate	1,000	mg
Vitamin B6	200	mg
Biotin	14,000	mcg
Folic Acid	100	mg
Vitamin B12	800	mcg
Choline Chloride	24,000	mg
Manganese	4,800	mg
Iron	800	mg
Zinc	4,800	mg
Copper	640	mg
Iodine	50	mg
Selenium	12	mg
DL-Methionine	10.8	%
L-Lysine	9.61	%
L-Threonine	7.12	%
Phosphor	8.535	%
Phosphor (AV)	15.335	%
Sodium	6.46	%
Chloride	8.52	%
6-Fytaze	80,000	FTU
Xylanase	60,000	FTU
Anti-Toxin	Included	
Anti-Oxidant	Included	
Anti-Caking	Included	



### DOSAGE:

25 Kg for 1 Ton of Feed.





25 kg

IT'S  
USED FOR  
CHICKENS



## ADEMIX BROILER GROWER 2.5%



### COMPOSITIONS

Each 1 kg contains:


Vitamin A	440,000	IU
Vitamin D3	180,000	IU
Vitamin E	2,600	mg
Vitamin K3	144	mg
Vitamin B1	160	mg
Vitamin B2	320	mg
Niacin	2,600	mg
Calcium D-Pantothenate	800	mg
Vitamin B6	160	mg
Biotin	11,200	mcg
Folic Acid	80	mg
Vitamin B12	720	mcg
Choline Chloride	24,000	mg
Manganese	4,800	mg
Iron	800	mg
Zinc	4,800	mg
Copper	640	mg
Iodine	50	mg
Selenium	12	mg
DL-Methionine	9.74	%
L-Lysine	6.98	%
L-Threonine	5.2	%
Phosphor	5.45	%
Phosphor (AV)	12.25	%
Sodium	6.71	%
Chloride	8.56	%
6-Fytaze	80,000	FTU
Xylanase	60,000	FTU
Anti-Toxin	Included	
Anti-Oxidant	Included	
Anti-Caking	Included	



### DOSAGE:

25 Kg for 1 Ton of Feed.





25 kg

IT'S  
USED FOR  
CHICKENS



## ADEMIX BROILER FINISHER 2.5%



### COMPOSITIONS


Each 1 kg contains:

Vitamin A	400,000	IU
Vitamin D3	160,000	IU
Vitamin E	2,200	mg
Vitamin K3	128	mg
Vitamin B1	120	mg
Vitamin B2	280	mg
Niacin	2,000	mg
Calcium D-Pantothenate	600	mg
Vitamin B6	120	mg
Biotin	8,800	mcg
Folic Acid	72	mg
Vitamin B12	640	mcg
Choline Chloride	24,000	mg
Manganese	4,800	mg
Iron	800	mg
Zinc	4,800	mg
Copper	640	mg
Iodine	50	mg
Selenium	12	mg
DL-Methionine	9.74	%
L-Lysine	6.98	%
L-Threonine	5.2	%
Phosphor	5.45	%
Phosphor (AV)	12.25	%
Sodium	6.71	%
Chloride	8.56	%
6-Fytaze	80,000	FTU
Xylanase	60,000	FTU
Anti-Toxin	Included	
Anti-Oxidant	Included	
Anti-Caking	Included	



### DOSAGE:

25 Kg for 1 Ton of Feed.



25 kg

 **IT'S  
USED FOR  
CHICKENS**



### DOSAGE:

- 0,5 - 0,75 Kg for 1 Ton of Feed.



**NOTE:** It Is Not Recommended To Be  
Used With Antibiotics

## PROBIOFEED



### COMPOSITIONS

Each 1 kg contains:

Lactobacillus Casei	60 x 10 <sup>8</sup>	CFU
Lactobacillus Acidophilus	240 x 10 <sup>8</sup>	CFU
Lactobacillus Plantarum	272 x 10 <sup>8</sup>	CFU
Lactobacillus Paracasei	392 x 10 <sup>8</sup>	CFU
Xylanase Endo-1.4 Beta Xylanase	1,872,000	IU
Beta Glucanase	294,000	IU
Amylase Alpha Amylase	168,000	IU
Protease	294,000	IU



### PROBIOFEED: WHAT IT DOES, HOW TO USE IT:

- Probiotics are beneficial bacteria and yeasts that help various body systems, especially the digestive system and skin, to work in a balanced way, unlike bacteria that cause infections and diseases.
- Probiotics, especially probiotics in the gut, are largely aimed at ensuring the proper functioning of the digestive and immune systems. Beneficial bacteria play a major role in maintaining the intestinal flora balance. Endikasyonları:

#### FOR ALL POULTRY

1. Contributes to the support of the immune system.
2. It helps digest food.
3. It helps the production of vitamins synthesized in the body.
4. It helps the cells that make up the intestinal wall and prevent harmful bacteria entering the body from reaching the blood.
5. It increases the absorption of drugs taken from the outside into the body.
6. Helps maintain the physiological balance in the intestine.
7. Eliminates bloating problems.
8. Protects liver and bone health.
9. Controls endemic and zoonotic agents by competitive exclusion.
10. It slows down the infectious agents that cause deadly diseases (such as E.coli, Salmonella, Campylobacter).
11. It provides an increase in egg and fertility.



# RUMINANTS PREMIXES

SUPER MILK SC

MILKMAX-SC

ADEMILK SC

CALF VIT

SUPER BEEF SC

BEEFMAX-SC

KOYUVIT - SC

MULTI MIX FORT

ADEVIT SUPER

MINERAMIX

SUPER TAMPOM

ADE - TAMPON

TAMPOMIX

SACC MIX








# OUR PREMIXES PRODUCTS

## RUMINANTS

Why are feed additives needed to provide extra nutrition?

It has been known for a long time that general raw feed composed of basic raw materials such as soybeans, corn and fat does not contain all the necessary nutrients, which makes the feed insufficient for the animal to obtain the best performance. Another fact is that problems such as low digestibility, microbial contamination, and tarnished color can be fixed by adding ingredients such as vitamins, enzymes, and amino acids are added to livestock feed. These two facts above are just two normal examples where feed additives are useful and helpful. To speak in a sum way, feed additives are the key to help animals to get complete nutrition and therefore have a healthier body condition. Since the relationship between the health of humans and the health of animals is very close, helping animals to become the best of themselves is helpful for humans as well. The main reason why feed additives are used so often in farm industries nowadays is that they are equipped with a great variety of nutrition. These special nutritions not only help to give animals the health they want but also help to maximize the nutrition of feedstuff for animals, thereby better the production efficiency in farm industries.





25 kg

IT'S USED FOR:



**BYPASS  
FAT**

# SUPER MILK SC



## COMPOSITIONS

Each 10 kg contains:

Vitamin A	14,500,000	IU
Vitamin D3	2,700,000	IU
Vitamin E	52,000	mg
Niacin	80,000	mg
Biotin	2,000	mg
DL-Methionine	2,000	mg
L-Lysine	2,000	mg
Manganese	40,000	mg
Manganese (Manganese Glysinate)	10,000	mg
Iron	40,000	mg
Zinc	40,000	mg
Zinc (Zinc Glysinate)	10,000	mg
Copper	8,000	mg
Copper (Copper Glysinate)	2,000	mg
Iodine	800	mg
Cobalt	300	mg
Selenium	200	mg
Phosphor	180,000	mg
Antioxidants (BHT)	10,000	mg
Saccharomyces Cerevisiae	100×10 <sup>10</sup>	CFU
Bypass Fat	1,300,000	mg
Toxin Binders	1,000,000	mg
Sodium Bicarbonate	100,000	mg
Vanilla Aroma	30,000	mg



## DOSAGE:

- 10 Kg for 1 Ton of Feed.
- 100 gr / day for calf and veal feed.
- 10 gr / day is added to cow and bull feed.



## INDICATIONS:

It is used to increase feed utilization of bovine and ovine animals. to ensure the development of young animals and to obtain a high level of efficiency. For milk breeds

- Increases feed utilization and increases appetite.
- Increases fertility and broking. Reveals hidden anger.
- Provides more meat and milk yield in animals.
- Increases peak time in lactation.
- It contributes to the acceleration of the development.
- Ensuring the healthy development of the offspring in pregnant animals and
- Reduces the likelihood of white muscle disease that may occur.
- It removes the disorders in the nails. Hair and skin.
- It reduces the risk of hypocalcemia (birth paralysis) that may occur after birth.
- Increases the animal's resistance against metabolic diseases.
- The fat ratio in milk increases.





25 kg

### IT'S USED FOR:



# MILKMAX-SC



## COMPOSITIONS

Each 10 kg contains:

Vitamin A	9,000,000	IU
Vitamin D3	2,000,000	IU
Vitamin E	30,000	mg
Niacin	60,000	mg
Biotin	750	mg
DL-Methionine	10,000	mg
L-Lysine	10,000	mg
Manganese	40,000	mg
Iron	50,000	mg
Zinc	40,000	mg
Copper	10,000	mg
Calcium	22,000	mg
Iodine	500	mg
Cobalt	200	mg
Selenium	200	mg
Magnesium Oxide	5,000	mg
Phosphor	5,000	mg
Sodium	3,000	mg
Saccharomyces Cerevisiae	100×10 <sup>10</sup>	CFU
Toxin Binders	2,000,000	mg
Antioxidants (BHT)	10,000	mg
Aroma	30,000	mg



## DOSAGE:

- 10 Kg for 1 Ton of Feed or
- 100 gr / day for calf and veal feed.
- 10 gr / day is added to cow and bull feed.



## INDICATIONS:

It is used to increase feed utilization of bovine and ovine animals. to ensure the development of young animals and to obtain a high level of efficiency. For milk breeds

- Increases feed utilization and increases appetite.
- Increases fertility and broking.
- Reveals hidden anger.
- Provides more meat and milk yield in animals.
- Increases peak time in lactation.
- It contributes to the acceleration of the development.
- Ensuring the healthy development of the offspring in pregnant animals and
- Reduces the likelihood of white muscle disease that may occur.
- It removes the disorders in the nails. Hair and skin.
- It reduces the risk of hypocalcemia (birth paralysis) that may occur after birth.
- Increases the animal's resistance against metabolic diseases.
- The fat ratio in milk increases.





25 kg

### IT'S USED FOR:



# ADEMILK SC



## COMPOSITIONS

Each 10 kg contains:

Vitamin A	6,000,000	IU
Vitamin D3	1,500,000	IU
Vitamin E	20,000	mg
Niacin	40,000	mg
Manganese	40,000	mg
Iron	40,000	mg
Zinc	40,000	mg
Copper	80,000	mg
Iodine	400	mg
Cobalt	100	mg
Selenium	100	mg
Saccharomyces Cerevisiae	100x10 <sup>10</sup>	CFU
DL-Methionine	5,500	mg
L-Lysine	4,400	mg
Magnesium Oxide	20,000	mg



## DOSAGE:

- 10 Kg for 1 Ton of Feed.
- 100 gr / day for calf and veal feed.
- 10 gr / day is added to cow and bull feed.



## INDICATIONS:

It is used to increase feed utilization of bovine and ovine animals. to ensure the development of young animals and to obtain a high level of efficiency. For milk breeds

- Increases feed utilization and increases appetite.
- Increases fertility and broking. Reveals hidden anger.
- Provides more meat and milk yield in animals.
- Increases peak time in lactation.
- It contributes to the acceleration of the development.
- Ensuring the healthy development of the offspring in pregnant animals and
- Reduces the likelihood of white muscle disease that may occur.
- It removes the disorders in the nails. Hair and skin.
- It reduces the risk of hypocalcemia (birth paralysis) that may occur after birth.
- Increases the animal's resistance against metabolic diseases.
- The fat ratio in milk increases.



25 kg

### IT'S USED FOR:



### DOSAGE:

Small calves: It is given daily by mixing 10-15 Gr. Of feed per calf. Or as veterinarian recommendations.

# CALF VIT



### COMPOSITIONS

Each 1 kg contains:

Vitamin A	2,000,000 IU	Zinc	5,000 mg
Vitamin D3	300,000 IU	Zinc Glycinate	2,000 mg
Vitamin E	2,500 mg	Iron	5,000 mg
Vitamin B1	800 mg	Cobalt	15 mg
Vitamin B2	400 mg	Iodine	80 mg
Niacin	2,000 mg	Selenium	15 mg
Calcium-D-Pantothenate	1,200 mg	Cholin Chloride 60%	9,600 mg
Vitamin B6	500 mg	Saccharomyces Cerevisiae	100 × 10 <sup>10</sup> CFU
Biotin	5,000 mcg	Pecti	10,000 mg
Vitamin B12	2,000 mcg	Kaolin	15,000 mg
Manganese	5,000 mg	BHT	1,000 mg
Manganese Glycinate	1,000 mg	Silica	1,000 mg
Copper Copper Glycinate	1,000 mg 500 mg		



### INDICATIONS:

#### -For dairy and fattening calves

- It is a mixture containing high-density vitamin supplement. Biotin. Niacin. Live yeast and organic minerals in order to meet the vitamin and trace mineral substance needs of calves.
- The most important factor determining the profitability of cattle enterprises is to raise healthy calves.
- CALF VIT is a formula enriched with live yeast along with vitamins. Organic and inorganic minerals that calves need.
- Increases the calf's Growth rate. And is effective in solving respiratory tract problems.
- It supports beneficial bacteria in the digestive system and increases feed efficiency.
- Reduces diarrhea and prevents toxicity.
- CALF - VIT which helps you prevent calf deaths. Also minimizes the impact of calves during stress periods.



# SUPER BEEF SC



25 kg

## IT'S USED FOR:



## INDICATIONS:

It is used to increase feed utilization of bovine and ovine animals, to ensure the development of young animals and to obtain a high level of efficiency. For breeding breeds:

- It increases feed utilization and increases appetite.
- Provides more et yield in animals.
- Removes deformities in nails, hair and skin.
- It increases the animal's resistance against metabolic diseases.



## COMPOSITIONS

Each 10 kg contains:

Vitamin A	8,000,000	IU
Vitamin D3	1,750,000	IU
Vitamin E	10,000	mg
Vitamin B1	1,960	mg
Vitamin B2	1,600	mg
Niacin	9,900	mg
Calcium-D-Pantothenate	3,000	mg
Vitamin B6	2,480	mg
Biotin	50	mg
Vitamin B12	10	mg
Copper	6,500	mg
Manganese	44,950	mg
Zinc	21,750	mg
Iron	39,000	mg
Iodine	680	mg
Selenium	50	mg
Cobalt	130	mg
Antioxidants (BHT)	10,000	mg
Saccharomyces Cerevisiae	300×10 <sup>10</sup>	CFU
Toxin Binders	525,000	mg
Aroma	30,000	mg




## DOSAGE:

- 10 Kg for 1 Ton of Feed.
- 50 gr / day for calf and veal feed.
- 100 gr / day is added to cow and bull feed.







25 kg

### IT'S USED FOR:



### DOSAGE:

- 10 Kg for 1 Ton of Feed.
- 50 gr / day for calf and veal feed.
- 100 gr / day is added to cow and bull feed.

# BEEFMAX-SC



### COMPOSITIONS

Each 10 kg contains:

Vitamin A	6,500,000	IU
Vitamin D3	500,000	IU
Vitamin E	6,000	mg
Biotin	50	mg
Iron	30,000	mg
Copper	3,000	mg
Zinc	30,000	mg
Manganese	40,000	mg
Iodine	600	mg
Cobalt	100	mg
Selenium	100	mg
Phosphor	9,000	mg
Saccharomyces Cerevisiae	100 × 10 <sup>10</sup>	CFU
Toxin Binders	50,000	mg
Antioxidants (BHT)	10,000	mg
Aroma	30,000	mg



### INDICATIONS:


It is used to increase feed utilization of bovine and ovine animals.

To ensure the development of young animals and to obtain a high level of efficiency.

For breeding breeds:

- It increases feed utilization and increases appetite.
- Provides more ET yield in animals.
- Removes deformities in nails, Hair and skin.
- It increases the animal's resistance against metabolic diseases.





25 kg

#### IT'S USED FOR:



#### DOSAGE:

- Sheep, Goats, Cattle milk feeds, Beef fattening feeds  
Lamb, Calf starter and Rearing 10 Kg is added to  
the ton of their feed.

## KOYUVIT – SC



#### COMPOSITIONS

Each 10 kg contains:

Vitamin A	7,500,000	IU
Vitamin D3	1,000,000	IU
Vitamin E	15,000	mg
Vitamin B1	1,960	mg
Vitamin B2	1,600	mg
Niacin	9,900	mg
Calcium-D-Pantothenate	3,000	mg
Vitamin B6	2,480	mg
Biotin	30	mg
Vitamin B12	10	mg
Copper	3,500	mg
Manganese	49,600	mg
Zinc	29,000	mg
Iron	30,000	mg
Iodine	810	mg
Selenium	900	mg
Cobalt	40	mg
Saccharomyces Cerevisiae	200×10 <sup>10</sup>	CFU
Toxin Binders	750,000	mg
Aroma	30,000	mg



#### INDICATIONS:

- Provides essential vitamins and trace minerals for cattle and sheep milk feeds.
- By increasing feed consumption. it helps them Grow healthier and faster.
- Increases resistance to diseases by strengthening the immune system.
- Increases milk yield and quality.
- Increases peak time in lactation.
- Increases the quality of carcass.
- It is protective against pica disease (tree. wall licking. bag eating).
- It reduces the risk of hypocalcemia (birth paralysis) that may occur after birth.







1000 gr

IT'S USED FOR:



# MULTI MIX FORT



## COMPOSITIONS

Each 1000 gr contains:

Vitamin A	6,000,000	IU
Vitamin D3	1,500,000	IU
Vitamin E	5,000	mg
Vitamin B1	1,000	mg
Vitamin B2	2,000	mg
Niacin	5,000	mg
Calcium-D-Pantothenate	2,000	mg
Vitamin B6	1,000	mg
Biotin	3,000	mcg
Folic Acid	250	mg
Vitamin B12	8,000	mcg
Manganese Oxide	10,000	mg
Copper Sulphate	3,000	mg
Zinc Sulphate	2,000	mg
Iron Sulphate	4,000	mg
Cobalt Sulphate	20	mg
Calcium Iodate	20	mg
Sodium Selenite 4.5%	5	mg
DL-Methionine	15,000	mg
Choline Chloride 60%	15,000	mg
Calcium Carbonate	Up To 1,000	gr



## INDICATIONS:

- Prevention or treatment of vitamin and Mineral deficiencies in farm animals.
- Improvement of feed transitions.
- Minerals play a key role in the maintenance of osmotic pressure, and thus regulate the exchange of water and solutes within the animal body.
- Minerals serve as structural constituents of soft tissues.
- Minerals are essential for the transmission of nerve impulses and muscle contraction.
- Minerals play a vital role in the acid-base equilibrium of the body, and thus regulate the pH of the blood and other body fluids.



## DOSAGE:

1 kg per 1 ton of feed.





# ADEVIT SUPER



## COMPOSITIONS

Each 1000 gr contains:

Vitamin A	4,000,000	IU
Vitamin D3	1,000,000	IU
Vitamin E	15,000	mg
Manganese Oxide	16,000	mg
Copper Sulphate	1,000	mg
Zinc Sulphate	5,000	mg
Iron Sulphate	8,000	mg
Cobalt Sulphate	40	mg
Calcium Iodate	100	mg
Sodium Selenite 4.5%	200	mg
Calcium Carbonate	Up To 1,000	gr



## INDICATIONS:

- Prevention or treatment of vitamin and Mineral deficiencies in farm animals.
- Improvement of feed transitions.
- Minerals play a key role in the maintenance of osmotic pressure, and thus regulate the exchange of water and solutes within the animal body.
- Minerals serve as structural constituents of soft tissues.
- Minerals are essential for the transmission of nerve impulses and muscle contraction.
- Minerals play a vital role in the acid-base equilibrium of the body, and thus regulate the pH of the blood and other body fluids.



## DOSAGE:

1 Kg per ton of feed .



1000 gr

## IT'S USED FOR:





1000 gr

### IT'S USED FOR:



# MINERAMIX



## COMPOSITIONS

Each 1000 gr contains:

Manganese Sulphate	4,000	mg
Copper Sulphate	60,000	mg
Zinc Oxide	2,000	mg
Iron Sulphate	1,000	mg
Magnesium Sulphate	80,000	mg
Cobalt Sulphate	20	mg
Sodium Chloride	125,000	mg
Phosphor	21,000	mg
Calcium	17,182	mg
Calcium Carbonate	Up To 1,000	gr



## INDICATIONS:

- Prevention or treatment of Mineral deficiencies in farm animals.
- Improvement of feed transitions.
- Minerals play a key role in the maintenance of osmotic pressure, and thus regulate the exchange of water and solutes within the animal body.
- Minerals serve as structural constituents of soft tissues.
- Minerals are essential for the transmission of nerve impulses and muscle contraction.
- Minerals play a vital role in the acid-base equilibrium of the body, and thus regulate the pH of the blood and other body fluids.



## DOSAGE:

1 Kg per ton of feed .





25 kg

IT'S USED FOR:



# SUPER TAMPOM



## COMPOSITIONS

Each 10 kg contains:

Sodium Bicarbonate	6,000,000	mg
Magnesium Oxide	2,000,000	mg
Potassium Chloride	68,000	mg
Sodium Chloride	132,000	mg
Inactive Yeast	175,000	mg



## INDICATIONS:

In capital animals:

- With the help of sodium bi carbonate in the content, the changes and deposits occurring in rumen's ph.
- In case of feed acidosis at tmrs with high concentrate feed ratio.
- In digestive problems.
- In relieving blasting and gaught occurring at the end of fattening.
- In hot stress.
- In case of acidosis after feeding and foot nail problems.
- Used to increase the use of feed.



## DOSAGE:

- Cattle: 100 gr (daily).
- Sheep: 20 gr (daily).





# ADE – TAMPON



25 kg

## IT'S USED FOR:



## COMPOSITIONS

Each 10 kg contains:

Sodium Bicarbonate	3,000,000	mg
Magnesium Oxide	2,000,000	mg
Potassium Chloride	68,000	mg
Sodium Chloride	132,000	mg
Inactive Yeast	175,000	mg



## INDICATIONS:

In capital animals:

- With the help of sodium bi carbonate in the content. the changes and deposits occuring in rumen's ph.
- In case of feed acidosis at tmrs with high concentrate feed ratio.
- In digestive problems.
- In relieving blasting and gaught occuring at the end of fattening.
- In hot stress.
- In case of acidosis after feeding and foot nail problems.
- Used to increase the use of feed.



## DOSAGE:

- Cattle: 150 gr (daily).
- Sheep: 30 gr (daily).





25 kg

IT'S USED FOR:



# TAMPOMIX



## COMPOSITIONS

Each 10 kg contains:

Sodium Bicarbonate	2,250,000	mg
Magnesium Oxide	1,100,000	mg
Potassium Chloride	25,000	mg
Sodium Chloride	75,000	mg
Inactive Yeast	175,000	mg



## INDICATIONS:

In capital animals:

- With the help of sodium bi carbonate in the content. the changes and deposits occuring in rumen's ph.
- In case of feed acidosis at tmrs with high concentrate feed ratio.
- In digestive problems.
- In relieving blasting and gaught occuring at the end of fattening.
- In hot stress.
- In case of acidosis after feeding and foot nail problems.
- Used to increase the use of feed.



## DOSAGE:

- Cattle: 200 gr (daily).
- Sheep: 40 gr (daily).







25 kg

IT'S USED FOR:



# SACC MIX



## COMPOSITIONS

Each 1 kg contains:

Saccharomyces Cerevisiae 100 x10<sup>10</sup> CFU



## INDICATIONS:

- For Dairy and Livestock It provides maximum utilization of nutrients by regulating the rumen gut and flora.
- It increases the fattening performance of the male animal and the milk performance of the female animal.
- It regulates the stomach flora and helps to prevent gas problem.
- It provides live weight gain and quality milk production.
- Eliminates diarrhea that may occur during transition periods.
- It supports the development of bone, teeth and cartilage.
- It regulates and stabilizes the rumen microflora in regular and continuous use.



## DOSAGE:

1 Kg for 1 Ton of Feed.







Scan The QR Code  
For More Information



| ADEMIX VET

**Ademix yem sanayi ve ticaret limited řirketi.**

Adress: Çilek Mah. 63147 Sok, No:1/31 Akdeniz / Mersin - Türkiye

Website: [www.ademixvet.com](http://www.ademixvet.com) | Email: [info@ademixvet.com](mailto:info@ademixvet.com)

Tel: +90 533 073 45 97

Tel: +90 530 810 45 97